

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification 5 :</b> <b>C07H 21/02, 21/04, C12N 15/70</b> <b>C12N 15/74, 15/79, 5/10</b> <b>C07K 15/28, A61K 39/395</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 94/05690</b> <b>(43) International Publication Date:</b> 17 March 1994 (17.03.94)
<b>(21) International Application Number:</b> PCT/US93/08435 <b>(22) International Filing Date:</b> 8 September 1993 (08.09.93)  <b>(30) Priority data:</b> 07/941,654 9 September 1992 (09.09.92) US  <b>(60) Parent Application or Grant</b> (63) Related by Continuation US 07/941,654 (CIP) Filed on 9 September 1992 (09.09.92)  <b>(71) Applicants (for all designated States except US):</b> SMITH-KLINE BEECHAM CORPORATION [US/US]; 709 Swedeland Road, King of Prussia, PA 19406-2799 (US). UNITED STATES OF AMERICA as represented by THE SECRETARY OF THE NAVY [US/US]; Pentagon, Washington, DC 20031 (US). UNITED STATES OF AMERICA as represented by THE SECRETARY OF THE ARMY [US/US]; Pentagon, Washington, DC 20031 (US).		<b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only) :</b> GROSS, Mitchell Stuart [US/US]; 667 Pugh Road, Wayne, PA 19087 (US). ROSENBERG, Martin [US/US]; 241 Mingo Road, Royersford, PA 19468 (US). SADOFF, Jerald, Charles [US/US]; 1622 Kalmia Road, Washington, DC 20012 (US). HOFFMAN, Stephen [US/US]; 308 Argosy Drive, Gaithersburg, MD 20878 (US). SYLVESTER, Daniel, R. [US/US]; 42 Rossiter Avenue, Phoenixville, PA 19460 (US). CHAROENVIT, Yupin [US/US]; 2833 Schoolhouse Circle, Silver Spring, MD 20902 (US). HURLE, Mark, [US/US]; 641 Columbus Street, Bridgeport, PA 19405 (US).  <b>(74) Agents:</b> BAK, Mary, E. et al.; Howson and Howson, Spring House Corporate Center, P.O. Box 457, Spring House, PA 19477 (US).  <b>(81) Designated States:</b> AU, CA, JP, KR, NZ, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> NOVEL ANTIBODIES FOR CONFERRING PASSIVE IMMUNITY AGAINST INFECTION BY A PATHOGEN IN HUMANS  <b>(57) Abstract</b>  Proteins and peptides derived from a murine <i>P. falciparum</i> monoclonal antibody, including synthetic humanized variable light chain and variable heavy chain sequences, CDR peptides, and humanized antibodies useful in therapeutic methods and compositions for conferring passive immunity to infection by a malaria-causing parasite are provided.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NE	Niger
BE	Belgium	GN	Guinea	NL	Netherlands
BF	Burkina Faso	GR	Greece	NO	Norway
BG	Bulgaria	HU	Hungary	NZ	New Zealand
BJ	Benin	IE	Ireland	PL	Poland
BR	Brazil	IT	Italy	PT	Portugal
BY	Belarus	JP	Japan	RO	Romania
CA	Canada	KP	Democratic People's Republic of Korea	RU	Russian Federation
CF	Central African Republic	KR	Republic of Korea	SD	Sudan
CG	Congo	KZ	Kazakhstan	SE	Sweden
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovak Republic
CM	Cameroon	LU	Luxembourg	SN	Senegal
CN	China	LV	Latvia	TD	Chad
CS	Czechoslovakia	MC	Monaco	TG	Togo
CZ	Czech Republic	MG	Madagascar	UA	Ukraine
DE	Germany	ML	Mali	US	United States of America
DK	Denmark	MN	Mongolia	UZ	Uzbekistan
ES	Spain			VN	Viet Nam
FI	Finland				

NOVEL ANTIBODIES FOR CONFERRING PASSIVE IMMUNITY  
AGAINST INFECTION BY A PATHOGEN IN HUMANS

Field of the Invention

5 This invention relates generally to the field of monoclonal and recombinant antibodies directed to epitopes on selected pathogens, e.g., a malaria parasite, methods for preparing and using, and compositions employing, these antibodies.

Background of the Invention

10 Malaria is a severe and widespread disease, caused by various species of the protozoan parasite genus *Plasmodium*, including four species that infect man, e.g., *P. falciparum*, *P. vivax*, *P. ovale* and *P. malariae* [See, e.g., V. Enea et al., Science, 225:628-630 (1984)].  
15 Malaria remains one of the most widespread and fatal diseases in the world today because of the lack of an effective vaccine and programs to control vector populations, as well as new drug-resistant strains. Generally, treatment of malaria relies heavily on  
20 prophylactic drugs, such as the 4-aminoquinolines. However, for most cases, drug resistance by *P. falciparum* and the production of some undesirable side effects, have undermined the efficacy of these drug therapies.

The focus of much research effort in the field  
25 of malaria prophylaxis is the sporozoite form of the *Plasmodium* parasite, particularly the circumsporozoite (CS) protein [Clyde et al., Am. J. Trop. Med. Hyg., 24:397 (1975); Rieckman et al., Bull. WHO, 57(1):261 (1979); and U. S. Patent 4,957,869]. The cloning and  
30 characterization of the CS protein genes or fragments thereof of a number of *Plasmodium* species and recombinant expression thereof in *E. coli* or yeast host cells have been reported. The central repeat domain of the CS proteins is immunodominant, i.e., if one injects

sporozoites into an animal, the animal produces anti-repeat antibodies. The first anti-sporozoite candidate vaccine tested in man was based upon the repetitive epitopes found on the CS protein of *P. falciparum* consisting of (AsnAlaAsnPro)<sub>37</sub>(AsnValAspPro)<sub>4</sub> [SEQ ID NO: 1], which is invariant in a number of strains examined to date. Clinical trials utilizing a vaccine candidate, called R32tet32, consisting of NH<sub>2</sub>-Met-Asp-Pro-[(Asn-Ala-Asn-Pro)<sub>15</sub>(Asn-Val-Asp-Pro)<sub>1</sub>]<sub>2</sub>-Leu-Arg-Arg-Thr-His-Arg-Gly-Arg-His-His-Arg-Arg-His-Arg-Cys-Gly-Cys-Trp-Arg-Leu-Tyr-Arg-Arg-His-His-Arg-Trp-Gly-Arg-Ser-Gly-Ser-COOH [SEQ ID NO: 2], produced a protective response in a human volunteer against the sporozoite challenge [see, Ballou et al., The Lancet, June 6, 1987, pp. 1277-1281; and European Patent Publication No. 0192626, published August 27, 1986, incorporated herein by reference].

Numerous monoclonal antibodies (mAbs) directed toward proteins from various stages of the *Plasmodium* life cycle have been identified and shown to be effective in passive transfer experiments in mice and monkeys [Y. Charoenvit et al., J. Immunol., 146(3):1020-1025 (1990)]. However, the use of antibodies for the treatment or prophylaxis of malaria may have disadvantages. The administration of murine or other animal antibodies to humans may be limited by the adverse immune response of humans to the foreign antibody, e.g., rapid clearance and toxic side effects. Such immune responses in humans have been shown to be directed against both immunoglobulin constant and variable regions of murine antibodies.

Several techniques have been described which suggest alteration of murine (and other species) antibodies to reduce the occurrence of an immune response in a desired species, e.g., human, to the parent antibody [See, e.g., PCT Patent Publication No. PCT/WO86/01533, published March 13, 1986; British Patent Application No.

GB2188638A, published October 7, 1987; Amit et al., Science, 233:747-753 (1986); Queen et al., Proc. Natl. Acad. Sci. USA, 86:10029-10033 (1989); PCT Patent Publication No. PCT/WO90/07861, published July 26, 1990; and Riechmann et al., Nature, 332:323-327 (1988)]. While the prior art suggests possible experimental techniques, none show how to provide the combination of properties required for effective prevention of *in vivo* growth of *P. falciparum*.

There remains a need in the art for alternative methods of providing immunity against infection with selected pathogens, e.g., a malarial parasite, particularly for a prophylactic agent capable of providing effective short-term protection.

#### Summary of the Invention

In one aspect, the present invention provides complementarity determining region (CDR) peptides from a monoclonal antibody directed against a selected epitope on a pathogen, as well as fragments and analogs of these peptides. Preferably, the antibody is capable of binding an epitope of *Plasmodium*, particularly the CS repeat region epitope or a fragment thereof, e.g., murine anti-*P. falciparum* mAb NFS2. These CDRs retain the antigen binding specificity of the mAb from which they were derived.

Another aspect provides an isolated, naturally occurring or synthetic, humanized immunoglobulin light or heavy chain variable region amino acid sequence comprising one or more CDR sequences originating from the light or heavy chain of such a selected antibody.

In yet a further aspect, the invention provides a fusion protein comprising a first amino acid sequence derived from the variable light chain and/or heavy chain of an anti-*Plasmodium* antibody, an anti-*Plasmodium* CDR, a

functional fragment or analog thereof. The first selected amino acid sequence is operatively linked or fused to a second selected amino acid sequence. These fusion proteins are characterized by the antigen binding specificity of the mAb from which the first selected amino acid sequence is derived.

A further aspect of the invention provides an engineered antibody with specificity for the selected *Plasmodium* epitope, e.g., *P. falciparum* repeat region.

In another aspect, the invention provides a *P. falciparum* antibody or fragment thereof produced by screening hybridoma products derived from any species immunoglobulin repertoires, or human or murine antibody combinatorial libraries, with the epitope of mAb NFS2.

In a further aspect, the present invention provides  $F_{ab}$  fragments of the above-described engineered antibodies or anti-*Plasmodium* mAbs.

As yet additional aspects, the invention provides nucleic acid sequences which encode the proteins, peptides, antibodies and fragments described herein, as well as plasmids containing one or more of the sequences, host cells transformed therewith, and methods for producing the products of expression of these nucleotide sequences in host cells, e.g., mammalian cells.

Other aspects provided by the invention include a pharmaceutical composition and a prophylactic method for conferring passive immunity to a human anticipating exposure to a malarial parasite, comprising an effective amount of at least one protein, antibody, peptide or fragment described herein and a pharmaceutically acceptable carrier or diluent.

Other aspects and advantages of the present invention are described further in the following detailed description of preferred embodiments thereof.

Brief Description of the Drawings

Fig. 1 illustrates the amino acid SEQ ID NO: 4 and nucleotide SEQ ID NO: 3 sequences of the naturally occurring light chain variable region of mAb NFS2.

Fig. 2 illustrates the amino acid SEQ ID NO: 6 and nucleotide SEQ ID NO: 5 sequences of a synthetic humanized light chain variable region Pfhzlc1-1 containing anti-*Plasmodium* CDRs SEQ ID NOs: 21-26. The CDRs are underlined.

Fig. 3 illustrates the amino acid SEQ ID NO: 8 and nucleotide SEQ ID NO: 7 sequences of synthetic humanized light chain variable region Pfhzlc1-2.

Fig. 4 illustrates the amino acid SEQ ID NO: 10 and nucleotide SEQ ID NO: 9 sequences of the naturally occurring heavy chain variable region of mAb NFS2.

Fig. 5 illustrates the amino acid SEQ ID NO: 12 and nucleotide SEQ ID NO: 11 sequences of a synthetic humanized heavy chain variable region Pfhzhc2-4.

Fig. 6 illustrates the amino acid SEQ ID NO: 14 and nucleotide SEQ ID NO: 13 sequences of synthetic humanized heavy chain variable region Pfhzhc2-3.

Fig. 7 is a schematic drawing of plasmid Pfhzhc2-3-Pcd employed to express a synthetic anti-*Plasmodium* heavy chain in mammalian cells. The plasmid contains a beta lactamase (Beta-lac) gene, an SV40 origin of replication (SV40), a cytomegalovirus promoter sequence (CMV), the synthetic heavy chain Pfhzhc2-3 SEQ ID NO: 13, a poly A signal from bovine growth hormone (BGH), a betaglobin promoter (beta glopro), a dihydrofolate reductase gene (DHFR), and another BGH sequence poly A signal in a pUC19 background.

Fig. 8 is a schematic drawing of plasmid Pfhzlc1-1-Pcn employed to express a synthetic light chain in mammalian cells. The plasmid differs from that of Fig. 7, in that it contains the synthetic humanized light

chain Pfhz1c1-1 SEQ ID NO: 5 rather than the heavy chain, and a neomycin gene (Neo) in place of DHFR.

Fig. 9 illustrates the nucleotide SEQ ID NO: 42 and amino acid SEQ ID NO: 43 sequences of a synthetic humanized heavy chain variable region Pfzhzc2-6.

#### Detailed Description of the Invention

The present invention provides prophylactic agents capable of conferring a short duration, protective immune state against infection of humans by selected pathogens in the immunized human, e.g., for epidemic control and for use by those anticipating exposure to the pathogen. Recombinant or engineered antibodies, preferably chimeric, humanized or human monoclonal antibodies, are capable of use as such passive protective proteins. These proteins in a prophylactic composition may be administered before anticipated exposure to the pathogen and would not require daily regimens of follow-up doses to mediate the short term protection.

While the following description refers specifically to antibodies capable of conferring passive protection to the sporozoite form of the pathogen, *P. falciparum*, a causative agent of malaria in humans, the invention described herein is not limited to any particular stage of that pathogen nor to that pathogen alone. The teachings of the present invention permit one skilled in the art to construct other recombinant antibodies directed to other selected pathogens, e.g., other species of *Plasmodium*, including the blood stages, liver stages, or gametocyte stages. Antibodies of the invention directed against the circumsporozoite CS gene of the other human infective parasites, e.g., *P. malariae*, *P. vivax* and *P. ovale*, may also be constructed according to this invention to provide passive transfer proteins useful against these parasitic infections.



Similarly, passive therapy agents prepared according to the invention may involve other infective agents, viruses, bacteria and the like. Additionally, such antibodies may also be useful as therapeutic agents for the treatment of acute stages of infections.

#### I. Definitions

"First fusion partner" refers to a nucleic acid sequence encoding an amino acid sequence, which can be all or part of an immunoglobulin heavy chain, a light chain, functional fragment thereof including the variable region from one or both chains and CDRs therefor, or an analog thereof, having the antigen binding specificity of a selected high titer antibody, preferably the murine antibody, NFS2.

"Second fusion partner" refers to another nucleotide sequence encoding a protein or peptide to which the first fusion partner is fused in frame or by means of an optional conventional linker sequence. Such second fusion partner is preferably heterologous to the first fusion partner. A second fusion partner may include a nucleic acid sequence encoding a second antibody region of interest, e.g., all or part of an appropriate human constant region or framework region.

"Fusion molecule" refers to the product of a first fusion partner operatively linked to a second fusion partner. "Operative linkage" of the fusion partners is defined as an association which permits expression of the antigen specificity of the anti-*P. falciparum* sequence (the first fusion partner) from the donor antibody as well as the desired characteristics of the second fusion partner. For example, a nucleic acid sequence encoding an amino acid linker may be optionally used, or linkage may be via fusion in frame to the second fusion partner.

"Fusion protein" refers to the protein encoded by the fusion molecule, which may be obtained by expression of the fusion molecule in a selected host cell. Such fusion proteins may be engineered antibodies, e.g., chimeric or humanized antibodies, or any of the antibody regions identified herein fused to immunoglobulin or non-immunoglobulin proteins and the like.

"Donor antibody" refers to an antibody (polyclonal, monoclonal or recombinant) which contributes its naturally-occurring or modified variable light and/or heavy chains, variable regions thereof, CDRs thereof or other functional fragments thereof to a first fusion partner, so as to provide the fusion molecule and fusion protein, with the antigenic specificity characteristic of the donor antibody. One donor antibody suitable for use in this invention is murine mAb NFS2 and others are described below.

"Acceptor antibody" refers to an antibody (polyclonal, monoclonal or recombinant) heterologous to the donor antibody, but homologous to the patient (human or other mammal) to be treated, which contributes all or any portion of the sequences of its variable heavy and/or light chain framework regions and/or its heavy and/or light chain constant regions to a second fusion partner. Preferably a human antibody is the acceptor antibody.

"CDRs" are defined as the complementarity determining region amino acid sequences of an antibody which are the hypervariable regions of the heavy and light chains. CDRs provide the majority of contact residues for the binding of the antibody to the antigen or epitope. CDRs of interest in this invention are derived from donor antibody variable heavy and light chain sequences, and include functional fragments and analogs of the naturally occurring CDRs, which share or

retain the same antigen binding specificity as the donor antibody from which they were derived.

By "sharing the antigen binding specificity" it is meant, for example, that although mAb NFS2 may be characterized by a certain level of antigen affinity, and a CDR encoded by a nucleic acid sequence of NFS2 in an appropriate structural environment may have a lower affinity, it is expected that CDRs of NFS2 in such environments will nevertheless recognize the same epitope(s) as NFS2.

A "functional fragment" is a partial CDR sequence or partial heavy or light chain variable sequence which retains the same antigen binding specificity as the antibody from which the fragment was derived.

An "analog" is an amino acid or peptide sequence modified by replacement of at least one amino acid, modification or chemical substitution of an amino acid, which modification permits the amino acid sequence to retain the biological characteristics, e.g., antigen specificity, of the unmodified sequence.

An "allelic variation or modification" is an alteration in the nucleic acid sequence encoding the amino acid or peptide sequences of the invention. Such variations or modifications may be due to degeneracies in the genetic code or may be deliberately engineered to provide desired characteristics. These variations or modifications may or may not result in alterations in any encoded amino acid sequence.

An "engineered antibody" is a type of fusion protein, i.e., a synthetic antibody (e.g., a chimeric or humanized antibody) in which a portion of the light and/or heavy chain variable domains of a selected acceptor antibody is replaced by analogous parts of CDRs from one or more donor antibodies which have specificity

for the selected epitope. These engineered antibodies may also be characterized by alteration of the nucleic acid sequences encoding the acceptor antibody light and/or heavy variable domain framework regions in order to retain donor antibody binding specificity. These antibodies can comprise immunoglobulin constant regions and variable framework regions from the acceptor antibody, and one or more CDRs from the *Plasmodium* donor antibodies described herein. Preferably the engineered antibodies of the invention will be produced by recombinant DNA technology.

"Chimeric antibody" refers to a type of engineered antibody which contains naturally-occurring variable region light chain and heavy chains (both CDR and framework regions) derived from a non-human donor mAb in association with light and heavy chain constant regions derived from a human (or other heterologous animal) acceptor mAb.

"Humanized antibody" refers to an engineered antibody having its CDRs and/or other portions of its light and/or heavy chain variable domain framework regions derived from a non-human donor immunoglobulin, the remaining immunoglobulin-derived parts of the molecule being derived from one or more human immunoglobulins. Such antibodies can also include engineered antibodies characterized by a humanized heavy chain associated with a donor or acceptor unmodified light chain or a chimeric light chain, or vice versa.

"Effector agents" refers to non-protein carrier molecules to which the fusion proteins, and/or natural or synthetic light or heavy chain of the donor antibody or other fragments of the donor antibody may be associated by conventional means. Such non-protein carriers can include conventional carriers used in the diagnostic field, e.g., polystyrene or other plastic beads, or other

non-protein substances useful in the medical field and safe for administration to humans and animals. Other effector agents may include a macrocycle, for chelating a heavy metal atom, or a toxin, such as ricin. Such effector agents are useful to increase the half-life of the anti-*Plasmodium* derived amino acid sequences or to add to its properties.

## II. *Anti-Plasmodium Antibodies*

For use in constructing the recombinant antibodies of this invention as it relates to malaria-causing pathogens, non-human species may be employed to generate a desirable donor antibody upon presentment with an antigen from a *Plasmodium* strain capable of infecting humans. Conventional hybridoma techniques are employed to provide a hybridoma cell line secreting a non-human mAb to the selected antigen. As one example, the murine mAb, NFS2, has been identified as a desirable antibody which may be employed for use in developing a chimeric or humanized antibody of this invention.

Murine IgG mAb NFS2 is characterized by an antigen binding specificity to the repeat region of the *P. falciparum* CS protein. In *in vitro* assays, it prevented invasion of sporozoites into human hepatocytes or hepatoma cells. Analogous antibodies in the mouse model have conferred passive protection against malaria. The production of mAb NFS2 is described in detail in Example 1 below.

This invention is not limited to the use of the illustrative NFS2 mAb or its hypervariable sequences. Wherever in the following description the donor mAb is identified as NFS2, this designation is made for simplicity of description only. Other anti-*Plasmodium* antibodies may be substituted therefor. Suitable antibodies include, for example, the murine mAb 2A10, which is directed against the CS repeat protein or other

mAbs described in R. A. Wirtz et al, Bull WHO, 65:39-45 (1987).

Antibodies produced in other animals protected by immunization with sporozoites or a protective epitope of a selected *Plasmodium* species may be similarly employed in this invention as a source of protective anti-*Plasmodium* sequences.

For example, the *P. falciparum* CS protein repeat region protein R32tet32 NH<sub>2</sub>-Met-Asp-Pro-[(Asn-Ala-Asn-Pro)<sub>15</sub>(Asn-Val-Asp-Pro)<sub>1</sub>]<sub>2</sub>-Leu-Arg-Arg-Thr-His-Arg-Gly-Arg-His-His-Arg-Arg-His-Arg-Cys-Gly-Cys-Trp-Arg-Leu-Tyr-Arg-Arg-His-His-Arg-Trp-Gly-Arg-Ser-Gly-Ser-COOH SEQ ID NO: 2 may be employed to elicit both human and murine mAbs with binding specificity therefor. This repeat region protein is a suitable target for screening for neutralizing antibodies useful in prophylactic agents against malarial infection.

Similarly, the epitope to which NFS2 is responsive, and analogs thereof, may be useful in the screening and development of additional *P. falciparum* antibodies, for use in the development of prophylactic compositions for short-term protection of humans against malaria. Other epitopes of interest include non-repetitive flanking region epitopes, other repeat domains or various liver, and blood and sexual stage epitopes of *Plasmodium* species. Knowledge of these epitopes enables one of skill in the art to define synthetic, and to identify naturally-occurring, peptides which would be suitable to confer passive or active immunity against *P. falciparum* or other *Plasmodium* species. This knowledge also permits the production of mAbs useful in the prophylaxis of malarial infection in humans.

For example, other *P. falciparum* antibodies may be developed by screening hybridomas or other combinatorial libraries, or antibody phage displays [W.

D. Huse et al., Science, 246:1275-1281 (1988)] using the murine mAb epitope described herein. A collection of antibodies, including hybridoma products or antibodies derived from any species immunoglobulin repertoire may be  
5 screened in a conventional competition assay, such as described in the examples below, with one or more epitopes described herein.

Antibodies such as those described above, including those generated against a desired epitope and  
10 produced by conventional techniques, including without limitation, genes encoding murine mAbs, human mAbs, and combinatorial antibodies, may be useful as donor antibodies, as sources of antibody fragments, as well as in prophylactic compositions against *P. falciparum* in  
15 humans. Preferably, the antibodies developed in response to *Plasmodium*, particularly *P. falciparum*, epitopes may be useful as donors of desirable variable heavy and/or light chain amino acid sequences, or functional fragments thereof (e.g., CDRs) useful in the development of fusion  
20 proteins, including engineered antibodies. Thus, the invention may utilize a donor antibody, other than NFS2, which is capable of binding to the *P. falciparum* peptide consisting essentially of the amino acid sequence of the repeat protein and analogs thereof.

25 Additionally, the mAbs identified herein, other mAbs which are developed and are responsive to the use of the sporozoites, R32tet32 [SEQ ID NO: 2] or the repeat epitopes identified herein may be further altered or manipulated to impart additional desirable prophylactic  
30 characteristics.

### III. Antibody Fragments, Amino Acid and Nucleotide Sequences

The present invention provides isolated naturally-occurring or synthetic variable light chain and  
35 variable heavy chain sequences derived from mAb NFS2, as

well as CDRs and fragments therefrom, which may be employed in the design of fusion proteins (including engineered antibodies) which are characterized by the antigen binding specificity of this mAb.

5           The naturally-occurring variable heavy chain of NFS2 is characterized by the amino acid and encoding nucleic acid sequences illustrated in Fig. 4 [SEQ ID NOS: 9 and 10]. This chain is characterized by CDRs having the following nucleotide and predicted amino acid  
10 sequences. CDR 1 is characterized by the sequence:

AGCTATGCCATGTCT SEQ ID NO: 32  
SerTyrAlaMetSer SEQ ID NO: 33.

The naturally occurring CDR 2 nucleic acid and amino acid sequences are SEQ ID NOS: 17 and 18, respectively:

15 GAAATTAGTGATGGTGGTAGTTACACCTACTATCCAGACACTGTGACGGGC  
GluIleSerAspGlyGlySerTyrThrTyrTyrProAspThrValThrGly.

The naturally-occurring CDR 3 has the nucleic acid and amino acid sequences SEQ ID NOS: 19 and 20, respectively:

20 CTCATCTACTATGGTTACGACGGGTATGCTATGGACTAC  
LeuIleTyrTyrGlyTyrAspGlyTyrAlaMetAspTyr.

Synthetic humanized variable heavy chains of NFS2 are characterized by the amino acid and encoding nucleic acid sequences illustrated in Fig. 5 [SEQ ID NOS: 9 and 10] and Fig. 6 [SEQ ID NOS: 13 and 14]. In both  
25 synthetic chains, the CDRs have the following nucleotide and predicted amino acid sequences. Nucleotide changes were made in CDR 1 from the naturally occurring CDRs, and are indicated by underlining. Synthetic CDR 1 is characterized by the sequence:

30 AGCTATGCCATGAGC SEQ ID NO: 15  
SerTyrAlaMetSer SEQ ID NO: 16.

The synthetic CDR 2 nucleic acid and amino acid sequences are identical to the naturally-occurring sequences SEQ ID NOS: 17 and 18, respectively. The synthetic CDR 3 has  
35 the same nucleic acid and amino acid sequences as does



15

the naturally occurring CDR 3 SEQ ID NOS: 19 and 20, respectively.

The naturally occurring variable light chain of NFS2 is characterized by the amino acid sequence and encoding nucleic acid sequence of Fig. 1 [SEQ ID NOS: 3 and 4]. This chain is further characterized by CDRs having the following nucleotide and amino acid sequences. CDR 1 is characterized by the nucleic acid and amino acid sequences SEQ ID NOS: 34 and 35, respectively:

5 AAGTCCAGTCAGAGCCTTTTATATAGTAGCAATCAAAGAATTACTTGGCC  
LysSerSerGlnSerLeuLeuTyrSerSerAsnGlnLysAsnTyrLeuAla.

CDR 2 is characterized by the nucleic acid and amino acid sequences SEQ ID NOS: 36 and 37, respectively:

10 TGGGCATCCACTAGGGAATCT  
15 TrpAlaSerThrArgGluSer.

CDR 3 is characterized by the nucleic acid and amino acid sequences SEQ ID NOS: 38 and 39, respectively:

CAGCAATATTATAGCTATCCTCGGACG  
GlnGlnTyrTyrSerTyrProArgThr.

20 A synthetic humanized variable light chain of NFS2 is characterized by the amino acid and encoding nucleic acid sequences illustrated in Fig. 2 [SEQ ID NOS: 5 and 6]. This chain is characterized by CDRs having the following predicted amino acid sequences and encoded by the illustrated nucleotide sequences. Nucleotide changes were made in the three CDRs from the naturally occurring corresponding CDRs, and are indicated by underlining.

Synthetic CDR 1 is characterized by the nucleic acid and amino acid sequences SEQ ID NOS: 21 and 22, respectively:

30 AAGAGCTCTCAGAGCCTTTTATACTCGAGCAATCAAAGAATTACTTGGCC  
LysSerSerGlnSerLeuLeuTyrSerSerAsnGlnLysAsnTyrLeuAla.

Synthetic CDR 2 is characterized by the nucleic acid and amino acid sequences SEQ ID NOS: 23 and 24, respectively:

35 TGGGCGTCAACTAGGGAATCT  
TrpAlaSerThrArgGluSer.

Synthetic CDR 3 is characterized by the amino acid sequences SEQ ID NO: 26 and encoding nucleotide sequences SEQ ID NO:25, respectively:

5 CAGCAATATTATAGCTATCCGCGGACG  
GlnGlnTyrTyrSerTyrProArgThr.

Another synthetic humanized variable light chain of NFS2 is characterized by the amino acid and encoding nucleic acid sequences illustrated in Fig. 3 [SEQ ID NOS: 7 and 8]. This chain is characterized by  
10 identical CDRs 1 and 3 as in the Fig. 2 synthetic sequences. However, a nucleotide change was made in the CDR 2 from both the naturally occurring corresponding CDR 2, and the Fig. 2 synthetic CDR 2. Double underlining is employed to indicate the change from the Fig. 2 synthetic  
15 sequences. Synthetic CDR 2 is characterized by the nucleic acid and amino acid sequences SEQ ID NOS: 40 and 41, respectively: TGGGCGTCGACTAGGGAATCT  
TrpAlaSerThrArgGluSer.

The present invention also includes the use of  
20 F<sub>ab</sub> fragments or F<sub>(ab')<sub>2</sub></sub> fragments. A F<sub>ab</sub> fragment contains the entire light chain and amino terminal portion of the heavy chain; and a F<sub>(ab')<sub>2</sub></sub> fragment is the fragment formed by two F<sub>ab</sub> fragments bound by disulfide bonds. MAb NFS2 and engineered antibodies derived  
25 therefrom and described below provide sources of F<sub>ab</sub> fragments and F<sub>(ab')<sub>2</sub></sub> fragments which can be obtained by conventional means, e.g. cleavage of the mAb with the appropriate proteolytic enzymes, papain and/or pepsin, or by recombinant methods. The F<sub>ab</sub> fragments or F<sub>(ab')<sub>2</sub></sub>  
30 fragments may be derived from any of the mAbs described above, as agents protective *in vivo* against infection by malarial pathogens, particularly *P. falciparum*.

The variable chain peptide sequences of murine mAb NFS2, its variable chain peptide sequences and CDRs,  
35 functional fragments, F<sub>ab</sub> fragments, and analogs thereof, and the nucleic acid sequences encoding them, may be

useful in obtaining various fusion molecules encoding  
desired fusion proteins, particularly engineered  
antibodies, and in methods for preparing and  
administering pharmaceutical compositions containing  
5 them.

The nucleic acid sequences of the invention, or  
fragments thereof, encoding the variable light chains and  
heavy chain peptide sequences or CDR peptides, or  
functional fragments thereof are used in unmodified form  
10 or are synthesized to introduce desirable modifications.  
The isolated naturally-occurring or synthetic nucleic  
acid sequences, which are derived from mAB NFS2 or from  
other desired anti-*Plasmodium* antibodies, may optionally  
contain restriction sites to facilitate insertion or  
15 ligation into a suitable nucleic acid sequence encoding a  
desired antibody framework region, ligation with  
mutagenized CDRs, or fusion with a nucleic acid sequence  
encoding a selected second fusion partner.

Taking into account the degeneracy of the  
20 genetic code, various coding sequences may be constructed  
which encode the variable heavy and light chain amino  
acid sequences, and CDR sequences of the invention, e.g.,  
Figs. 1 - 6 [SEQ ID NOS: 3-26], and functional fragments  
and analogs thereof which share the antigen specificity  
25 of the donor antibody. The isolated or synthetic nucleic  
acid sequences of this invention, or fragments thereof,  
encoding the variable chain peptide sequences or CDRs or  
functional fragments thereof can be used to produce  
fusion proteins, i.e. chimeric or humanized antibodies,  
30 or other engineered antibodies of this invention, when  
operatively combined with a second fusion partner.

These sequences are also useful for mutagenic  
insertion of specific changes within the nucleic acid  
sequences encoding the CDRs or framework regions, and for  
35 incorporation of the resulting modified or fusion nucleic

acid sequence into a vector for expression. For example, silent nucleotide substitutions may be made in the nucleotide sequences encoding the CDRs to create restriction enzyme sites to facilitate insertion of the mutagenic frameworks, or to modify the selected frameworks at nucleotide positions analogous to those of the donor antibody. Such mutations may include those inserted for the purpose of contributing to higher antigen binding affinity.

10           IV. *Fusion Molecules and Fusion Proteins*

Fusion molecules of this invention can encode engineered antibodies, chimeric antibodies and humanized antibodies. A desired fusion molecule may contain a first fusion partner encoding an amino acid sequence having the antigen specificity of a *Plasmodium* antibody directed against the amino acid sequence of the repeat protein and analogs thereof, operatively linked to a second fusion partner. Desirably the source of the first fusion partner is a selected mAb, e.,g., mAb NFS2, the source of nucleic acid sequences of Figs. 1 [SEQ ID NO: 3] and 4 [SEQ ID NO: 9].

A fusion molecule may encode an amino acid sequence for a naturally occurring variable heavy chain sequence of Fig. 4 [SEQ ID NOS: 9 and 10], a functional fragment or analog thereof, a naturally occurring variable light chain sequence of Fig. 1 [SEQ ID NO: 3 and 4], a functional fragment or analog thereof, or one or more NFS2 CDRs [SEQ ID NO: 15-26]. Another exemplary fusion molecule may encode a synthetic variable heavy and/or light chain from the donor mAb, such as those of Figs. 2 [SEQ ID NOS: 5 and 6], 3 [SEQ ID NOS: 7 and 8], 5 [SEQ ID NOS: 11 and 12], and 6 [SEQ ID NOS: 13 and 14], having the antigen specificity of *P. falciparum* antibody.

A desirable fusion molecule of this invention may be characterized by encoding an amino acid sequence

containing at least one, and preferably all of the CDRs [SEQ ID NOS: 15-26] of the variable region of the heavy and/or light chains of the murine antibody NFS2, or a functional fragment or analog thereof.

5           The second fusion partners are defined above, and may include a sequence encoding a peptide, protein or fragment thereof heterologous to the CDR-containing sequence having the antigen specificity of NFS2. One example is a sequence encoding a second antibody region  
10 of interest and may optionally include a linker sequence.

          The resulting fusion molecule may encode both anti-*P. falciparum* antigen specificity and the characteristic of the second fusion partner, e.g., a functional characteristic such as secretion from a  
15 recombinant host, or a therapeutic characteristic if the fusion partner itself encodes a therapeutic protein, or additional antigenic characteristics, if the fusion partner encodes a protein having its own antigen specificity.

20           If the second fusion partner is derived from another antibody, e.g., any isotype or class of immunoglobulin framework or constant region (preferably human), or the like, an engineered antibody is provided. Thus, for example, a fusion molecule of this invention  
25 may comprise a complete antibody molecule, having full length heavy and light chains (Figs. 4 [SEQ ID NOS: 9 and 10] and 1 [SEQ ID NOS: 3 and 4]). For example, the invention includes isolated naturally-occurring or synthetic nucleic acid sequences, which may encode  
30 variable region sequences, CDR peptides, fragments thereof derived from desired *Plasmodium* mAbs, any fragment of an engineered antibody, such as the  $F_{ab}$  or  $F_{(ab')_2}$  fragment, a heavy chain dimer, or any minimal recombinant fragment thereof such as an  $F_v$  or a single-  
35 chain antibody (SCA) or any other sequence encoding a

protein with the same specificity as the selected mAb, e.g., the *Plasmodium* mAb NFS2.

The first fusion partner may also be associated with effector agents as defined above, to which the first fusion partner may be operatively linked by conventional means, e.g., attached to the NFS2 encoding nucleic acids by a covalent bridging structure.

Fusion or linkage between the first fusion partners and the selected second fusion partner may be by way of any suitable means, e.g., by conventional covalent or ionic bonds, protein fusions, or heterobifunctional cross-linkers, e.g., carbodiimide, glutaraldehyde, and the like. Where the first fusion partner is associated with an effector agent, non-proteinaceous, conventional chemical linking agents may be used to fuse or join the anti-*P. falciparum* amino acid sequences to the effector agent. Such techniques are known in the art and readily described in conventional chemistry and biochemistry texts.

Additionally, conventional inert linker sequences which simply provide for a desired amount of space between the fusion partners or between the first fusion partner and the effector agent may also be constructed into the fusion molecule. The design of such linkers is well known to those of skill in the art.

Expression of such fusion molecules results in fusion proteins of this invention. One particularly desirable type of fusion protein includes the engineered antibody in which, at a minimum, fragments of the variable heavy and/or light domains of an acceptor mAb have been replaced by analogous parts of the variable light and/or heavy chains from one or more donor monoclonal antibodies, which include the *Plasmodium* mAbs described herein, such as NFS2.

One example of a particularly desirable engineered antibody is a humanized antibody, in which CDRs from a desired donor murine mAb are inserted into the framework regions of a human antibody. A preferred donor antibody is one directed against a *Plasmodium* epitope, preferably one specific for the repeat region epitope of *P. falciparum*. A particularly preferred donor antibody has all or a portion of the variable domain amino acid sequences of NFS2. In these humanized antibodies one, two or preferably three CDRs from the *Plasmodium* antibody heavy chain and/or light chain variable regions are inserted into the framework regions of a selected human antibody, replacing the native CDRs of that latter antibody.

Preferably, the variable domains in both human heavy and light chains have been altered by CDR replacement. This engineered humanized antibody thus preferably has the structure of a natural human antibody or a fragment thereof. Such humanized antibodies may or may not also include minimal alteration of the acceptor mAb light and/or heavy variable domain framework region in order to retain donor mAb binding specificity. The humanized antibody possesses the combination of properties required for effective prevention and treatment of infectious *P. falciparum* disease in animals or man.

The remainder of the engineered antibody may be derived from any suitable acceptor human immunoglobulin. A suitable human antibody may be one selected from a conventional database, e.g., the Kabat database, Los Alamos database, and Swiss Protein database, by homology to the nucleotide and amino acid sequences of the donor antibody. A human antibody characterized by a homology to the framework regions of the donor antibody (on an amino acid basis) may be suitable to provide a heavy

chain constant region and/or a heavy chain variable framework region for the insertion of the donor CDRs. A suitable acceptor antibody capable of donating light chain constant or variable framework regions may be selected in a similar manner. It should be noted that the acceptor antibody heavy and light chains are not required to originate from the same human antibody.

Desirably, the heterologous framework and constant regions are selected from the human immunoglobulin classes and isotypes, such as IgG (subtypes 1 through 4), IgM, IgA and IgE. However, the acceptor antibody need not comprise only human immunoglobulin protein sequences. For instance, a gene may be constructed in which a DNA sequence encoding part of a human immunoglobulin chain is fused to a DNA sequence encoding the amino acid sequence of a polypeptide effector or reporter molecule.

As one example, an engineered antibody may be encoded by a synthetic nucleic acid sequence encoding CDRs of the variable light chain region of NFS2 or a functional fragment thereof in place of at least a part of the nucleic acid sequence encoding the light chain variable region of an acceptor mAb, and a nucleic acid sequence encoding CDRs of the variable heavy chain region of NFS2 or a functional fragment thereof in place of at least a part of the nucleic acid sequence encoding the heavy chain variable region of an acceptor mAb, such as a human antibody. The resulting humanized antibody is characterized by the antigen binding specificity of mAb NFS2.

Alternatively, the engineered antibody (or the other monoclonal antibodies) of the invention may have attached to it an effector or reporter molecule. Alternatively, the procedure of recombinant DNA technology may be used to produce an engineered antibody



of the invention in which the  $F_c$  fragment or CH3 domain of a complete antibody molecule has been replaced by an enzyme or toxin molecule.

It will be understood by those skilled in the art that such an engineered antibody may be further modified by changes in variable domain amino acids without necessarily affecting the specificity of the donor antibody. It is anticipated that heavy and light chain amino acids may be substituted by other amino acids either in the variable domain frameworks or CDRs or both. Such engineered antibodies can be effective in prevention of productive malaria (e.g., by *P. falciparum*) infection in humans.

Additionally, the invention provides fusion proteins which are chimeric antibodies, as defined above. Such antibodies differ from the humanized antibodies described above by providing the entire donor antibody heavy chain and light chain variable regions, including framework regions, e.g., Figs. 1 [SEQ ID NOS: 3 and 4] and 4 [SEQ ID NOS: 9 and 10], fused to acceptor constant regions for both chains.

#### V. *Production of Proteins and Antibodies*

A fusion molecule, recombinant antibody or fusion protein of this invention is desirably constructed by recombinant DNA technology using genetic engineering techniques. The same or similar techniques may also be employed to generate other embodiments of this invention, e.g., to construct the chimeric or humanized antibodies, the synthetic light and heavy chains, the CDRs, and the nucleic acid sequences encoding them, as above mentioned.

A specific embodiment of the compositions of this invention is set out in Example 3 below using the CDRs of murine NFS2 and one or more selected human antibody light and heavy chain framework regions. Briefly described, a hybridoma producing the murine

antibody NFS2 is conventionally cloned, and the cDNA of its heavy and light chain variable regions is obtained by techniques known to one of skill in the art, e.g., the techniques described in Sambrook *et al.*, Molecular Cloning (A Laboratory Manual), 2nd edition, Cold Spring Harbor Laboratory (1989). The variable regions of the NFS2 are obtained using PCR primers, and the CDRs are identified using a known computer database, e.g, Kabat, for comparison to other antibodies.

Homologous framework regions of a heavy chain variable region from a human antibody were identified using the same databases, e.g., Kabat, and a human antibody having homology to NFS2 was selected as the acceptor antibody. The sequences of synthetic heavy chain variable regions containing the NFS2 CDRs within the human antibody frameworks were designed with optional nucleotide replacements in the framework regions for restriction sites. This designed sequence was synthesized by overlapping oligonucleotides, amplified by polymerase chain reaction (PCR), and corrected for errors.

A suitable light chain variable framework region was designed in a similar manner, resulting in two synthetic light chain variable sequences containing the NFS2 CDRs. See, Figs. 2 [SEQ ID NOS: 5 and 6] and 3 [SEQ ID NOS: 7 and 8]. As stated above, the source of the light chain is not a limiting factor of this invention.

These synthetic variable light and/or heavy chain sequences and the CDRs of mAb NFS2, and their encoding nucleic acid sequences, are employed in the construction of fusion proteins and engineered antibodies, preferably humanized antibodies, of this invention, by the following process. By conventional techniques, a DNA sequence is obtained which encodes the donor antibody variable heavy or light chain regions,

wherein at least the CDRs and those minimal portions of the acceptor mAb light and/or heavy variable domain framework region required in order to retain donor mAb binding specificity as well as the remaining  
5 immunoglobulin-derived parts of the antibody chain derived from a human immunoglobulin.

A conventional expression vector or recombinant plasmid is produced by placing these sequences encoding the fusion protein in operative association with  
10 conventional regulatory control sequences capable of controlling the replication and expression in, and/or secretion from, a host cell. Such regulatory sequences may be readily selected by one of skill in the art and are not intended as a limitation of the present  
15 invention. Regulatory sequence include promoter sequences, e.g., CMV promoter, and signal sequences which can be derived by one of skill in the art from antibodies.

A first vector can contain a sequence encoding  
20 a light chain-derived polypeptide. Similarly, a second expression vector is produced having a similar DNA sequence which encodes a complementary antibody light or heavy chain. Preferably at least the CDRs (and those minimal portions of the acceptor mAb light and/or heavy  
25 variable domain framework region required in order to retain donor mAb binding specificity) of the variable domain are derived from a donor antibody and the remaining immunoglobulin-derived parts of the antibody chain are derived from a human immunoglobulin in these  
30 vectors. Preferably this second expression vector is identical to the first, with the exception of the coding sequences and selectable markers, to ensure that each polypeptide chain is functionally expressed.

In another alternative, a single vector of the  
35 invention may be used, the vector including the sequence

encoding both light chain and heavy chain-derived polypeptides. The DNA in the coding sequences for the light and heavy chains may comprise cDNA or genomic DNA or both.

5           A selected host cell is co-transfected by conventional techniques with both the first and second vectors (or the single vector) to create the transfected host cell of the invention comprising both the recombinant or synthetic light and heavy chains. The  
10 transfected cell is then cultured by conventional techniques to produce the engineered antibody of the invention. The humanized antibody which includes the association of both the recombinant heavy chain and/or light chain is screened from culture by appropriate  
15 assay, such as an ELISA assay followed by the Inhibition of Sporozoite Invasion (ISI) assay described in the examples below. Similar conventional techniques may be employed to construct other fusion proteins of this invention.

20           Thus, the invention also includes a recombinant plasmid containing the coding sequence of the fusion molecule or engineered antibody of the invention. Such a vector is prepared by conventional techniques and suitably comprises the above-described DNA sequences and  
25 a suitable promoter operatively linked to the DNA sequences which encode the engineered antibody. Such a vector is transfected into a mammalian cell via conventional techniques.

          Suitable vectors for the cloning and subcloning  
30 steps employed in the methods and construction of the compositions of this invention may be selected by one of skill in the art. For example, the conventional pUC series of cloning vectors, may be used. One vector used is pUC19, which is commercially available from supply  
35 houses, such as Amersham (Buckinghamshire, United

Kingdom) or Pharmacia (Uppsala, Sweden). Additionally, any vector which is capable of replicating readily, has an abundance of cloning sites and marker genes, and is easily manipulated may be used for cloning. Thus, the selection of the cloning vector is not a limiting factor in this invention.

Similarly, the vectors employed for expression of the engineered antibodies according to this invention may be selected by one of skill in the art from any conventional vector. The vectors also contain selected regulatory sequences which are in operative association with the DNA coding sequences of the immunoglobulin regions and capable of directing the replication and expression of heterologous DNA sequences in selected host cells, such as CMV promoters. These vectors contain the above described DNA sequences which code for the engineered antibody or other fusion protein. Alternatively, the vectors may incorporate the selected immunoglobulin sequences modified by the insertion of desirable restriction sites for ready manipulation.

The expression vectors may also be characterized by marker genes suitable for amplifying expression of the heterologous DNA sequences, e.g., the mammalian dihydrofolate reductase gene (DHFR) or neomycin resistance gene (neo<sup>R</sup>). Other preferable vector sequences include a poly A signal sequence, such as from bovine growth hormone (BGH), and the betaglobin promoter sequence (betaglupro). The expression vectors useful herein may be synthesized by techniques well known to those skilled in this art.

The components of such vectors, e.g. replicons, selection genes, enhancers, promoters, and the like, may be obtained from natural sources or synthesized by known procedures for use in directing the expression of the recombinant DNA in a selected host. Other appropriate

expression vectors of which numerous types are known in the art for mammalian, bacterial, insect, yeast, and fungal expression may also be selected for this purpose.

Two exemplary expression vectors employed in the following examples for expression of the synthetic heavy and light chain sequences are the mammalian vectors Pfhzhc2-3-Pcd and Pfhzlc1-1-Pcn (see Figs. 7 and 8). However, this invention is not limited to the use of these illustrative pUC19-based vectors.

The present invention also encompasses a cell line transfected with a recombinant plasmid containing the coding sequences of the engineered antibodies or other fusion protein described by this invention. Host cells useful for the cloning and other manipulations of these cloning vectors are also conventional. However, most desirably, cells from various strains of *E. coli* are used for replication of the cloning vectors and other steps in the construction of the mAbs of this invention.

Suitable host cells or cell lines for the expression of the engineered antibody or other protein of the invention of this invention are preferably a eukaryotic cell, and most preferably a mammalian cell, such as a CHO cell or a myeloid cell. Other primate cells may be used as host cells and, most desirably, human cells are used, thus enabling the protein to be modified with human glycosylation patterns.

Alternatively, other eukaryotic cell lines may be employed. The selection of suitable mammalian host cells and methods for transformation, culture, amplification, screening and product production and purification are known in the art. See, e.g., Sambrook et al., cited above.

Bacterial cells may prove useful as host cells suitable for the expression of the recombinant mAbs of the present invention. However, due to the tendency of

proteins expressed in bacterial cells to be in an unfolded or improperly folded form or in a non-glycosylated form, any recombinant mAb produced in a bacterial cell would have to be screened for retention of antigen binding ability. If the protein expressed by the bacterial cell was produced in a properly folded form, that bacterial cell would be a desirable host. For example, various strains of *E. coli* used for expression are well-known as host cells in the field of biotechnology. Various strains of *B. subtilis*, *Streptomyces*, other bacilli and the like may also be employed in this method.

Where desired, strains of yeast cells known to those skilled in the art are also available as host cells, as well as insect cells and viral expression systems. See, e.g. Miller et al., Genetic Engineering, 8:277-298, Plenum Press (1986) and references cited therein.

The general methods by which the vectors of the invention may be constructed, transfection methods required to produce the host cells of the invention, and culture methods necessary to produce the fusion protein, and preferably an engineered antibody of the invention from such host cell are all conventional techniques. Likewise, once produced, the fusion proteins, preferably the engineered antibodies of the invention, may be purified from the cell culture contents according to standard procedures of the art, including ammonium sulfate precipitation, affinity columns, column chromatography, gel electrophoresis and the like. Such techniques are within the skill of the art and do not limit this invention.

The engineered antibody is then examined for *in vitro* activity by use of an assay appropriate for the selected pathogen. Presently conventional ELISA assay

formats are employed to assess qualitative and quantitative binding of the engineered antibody to the R32tet32 epitope [SEQ ID NO: 2]. The ISI assay described in Example 6 may also be employed. A similar assay, the inhibition of hepatocyte invasion assay (ILSDA), may be performed [S. Mellouk *et al.*, *Bull. WHO*, Suppl. 68:52-58 (1990)]. Additionally, assays currently being developed in the SCID mouse model may also be used to verify efficacy prior to subsequent human clinical studies performed to evaluate the persistence of the engineered antibody in the body despite the usual clearance mechanisms.

The examples below demonstrate the method for constructing the humanized antibodies derived from the murine mAb NFS2. Following the procedures described for humanized antibodies prepared from this antibody, one of skill in the art may also construct humanized antibodies from other malarial antibodies, variable region sequences and CDR peptides described herein. Engineered antibodies can be produced with variable region frameworks potentially recognized as "self" by recipients of the altered antibody. Minor modifications to the variable region frameworks can be implemented to effect large increases in antigen binding without appreciable increased immunogenicity for the recipient. Such engineered antibodies can effectively passively protect a human against *P. falciparum* infection.

#### VI. *Therapeutic/Prophylactic Uses*

The fusion proteins, particularly the engineered antibodies described above, functional fragments, analogs and the other protein or peptides described herein may be employed as prophylactic agents, capable of conferring short-term passive immunity to infection by the pathogen from which the original antigenic substance derives, e.g., *P. falciparum*, to a



subject. The protective effect conferred by the use of the engineered antibodies of this invention is produced by binding of the immunoglobulin to the pathogen and the subsequent removal of this bound complex by the normal function of macrophages. Thus, the engineered antibodies of the present invention, when in preparations and formulations appropriate for prophylactic use, are highly desirable for persons anticipating short-term exposure to the pathogen, e.g., travelers, tourists, or military personnel anticipating travel in endemic areas.

Therefore, this invention also relates to a method of prophylactic treatment of human *P. falciparum* infection in a human in need thereof which comprises administering an effective, protective dose of antibodies including one or more of the engineered antibodies or other fusion proteins described herein, or fragments thereof, to a human anticipating exposure to a species of *Plasmodium*.

The fusion proteins, including the engineered antibodies or fragments thereof of this invention, may also be used in conjunction with other antibodies, particularly human mAbs reactive with other epitopes responsible for the disease against which the engineered antibody of the invention is directed. Similarly mAbs reactive with other epitopes responsible for the disease in a selected animal against which the antibody of the invention is directed may also be employed in veterinary compositions. Any antibody that is capable of operating without interfering with the *Plasmodium* antibody of this invention, e.g., antibodies to other malaria stages or to different epitopes, are useful in these compositions.

The prophylactic agents of this invention are believed to be desirable to confer protection to exposure to the pathogen for from about 4 days to about 8 weeks, without requiring booster dosages of the agent. This

definition of 'short-term' relates to the relative duration of the recombinant antibodies of the present invention in the human circulation.

The mode of administration of the prophylactic agent of the invention may be any suitable route which delivers the agent to the host. The fusion proteins, including the engineered antibodies, and fragments thereof, and pharmaceutical compositions of the invention are particularly useful for parenteral administration, i.e., subcutaneously, intramuscularly or intravenously. However, the agent is preferably administered by intramuscular injection.

Prophylactic agents of the invention may be prepared as pharmaceutical compositions containing an effective amount of the engineered antibody of the invention as an active ingredient in a nontoxic and sterile pharmaceutically acceptable carrier. In the prophylactic agent of the invention, an aqueous suspension or solution containing the engineered antibody, preferably buffered at physiological pH, in a form ready for injection is preferred. The compositions for parenteral administration will commonly comprise a solution of the engineered antibody of the invention or a cocktail thereof dissolved in an acceptable carrier, preferably an aqueous carrier. A variety of aqueous carriers may be employed, e.g., saline, glycine, and the like. These solutions are sterile and generally free of particulate matter. These solutions may be sterilized by conventional, well known sterilization techniques. The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions such as pH adjusting and buffering agents, etc. The concentration of the antibody of the invention in such pharmaceutical formulation can

vary widely, i.e., from less than about 0.5%, usually at or at least about 1% to as much as 15 or 20% by weight and will be selected primarily based on fluid volumes, viscosities, etc., according to the particular mode of administration selected.

Thus, a pharmaceutical composition of the invention for parenteral, e.g., intramuscular injection, could be prepared to contain 1 mL sterile buffered water, and between about 50 to about 100 mg of an engineered antibody of the invention. Similarly, a pharmaceutical composition of the invention for intravenous infusion could be made up to contain 250 ml of sterile Ringer's solution, and 150 mg of an engineered antibody of the invention. Actual methods for preparing parenterally administrable compositions are well known or will be apparent to those skilled in the art and are described in more detail in, for example, Remington's Pharmaceutical Science, 15th ed., Mack Publishing Company, Easton, Pennsylvania.

It is preferred that the prophylactic agent of the invention, when in a pharmaceutical preparation, be present in unit dose forms. The appropriate therapeutically effective dose can be determined readily by those of skill in the art. To effectively prevent *P. falciparum* infection in a human or other animal, one dose of approximately 1 mg/kg to approximately 20 mg/kg of a protein or an antibody of this invention should be administered parenterally, preferably intramuscularly (i.m.) and possibly intravenously (i.v.). Such dose may be repeated at appropriate intervals during exposure.

The antibodies, engineered antibodies or fragments thereof described herein can be lyophilized for storage and reconstituted in a suitable carrier prior to use. This technique has been shown to be effective with

conventional immunoglobulins and art-known lyophilization and reconstitution techniques can be employed.

Single or multiple administrations of the pharmaceutical compositions can be carried out with dosage levels and pattern being selected by the treating physician. In any event, the pharmaceutical composition of the invention should provide a quantity of the engineered antibodies of the invention sufficient to effectively prevent infection.

The following examples illustrate the construction of exemplary engineered antibodies and expression thereof in suitable vectors and host cells, and are not to be construed as limiting the scope of this invention. All amino acids are identified by conventional codes or by full name, unless otherwise indicated. All restriction enzymes, plasmids, and other reagents and materials were obtained from commercial sources unless otherwise indicated. All general cloning, ligation and other recombinant DNA methodology were as performed in Sambrook et al., cited previously, or the first edition thereof.

#### Example 1 - Description of Production of NFS2

Murine IgG mAb NFS2 was made by repeated injection of *P. falciparum* sporozoites into mice followed by B cell fusion with a myeloma cell line. The murine mAb NFS2 is characterized by an antigen binding specificity to the repeat region of the *P. falciparum* CS protein. Specifically, the NFS2 mAb binds to the epitope, Pro Asn Ala Asn Pro Asn SEQ ID NO: 27. It is possible that the antibody also binds to a larger, or overlapping epitope on the repeat region.

This murine mAb, in *in vitro* assays, prevented invasion of sporozoites into human hepatocytes and hepatoma cells. Analogous antibodies in the mouse model have conferred passive protection against malaria and have been observed to be highly potent [see, e.g., R. A. Wirtz et al., Bull WHO, 65:39-45 (1987), incorporated herein by reference]. This antibody is available from the U. S. Naval Medical Research Institute.

### Example 2 - Cloning and Sequencing of NFS2

10                    Cytoplasmic RNA was prepared by the method of  
Favaloro et al., Meth. Enzymol., 65:718-749 (1980) from  
NFS2, and hybridoma cell lines. The following primers  
were used in the synthesis of Ig heavy ( $V_H$ ) and light  
( $V_L$ ) chain variable region cDNAs, respectively. The  $V_L$   
15 primers, #2580 and #2789, extended from HindII through  
XbaI and were made to conserved regions of murine RNA.

HindIII

20 #2580: 5'CCAGATGTAAGCTTCAGCTGACCCAGTCTCCA3' SEQ ID NO: 28  
PvuII

#2789:                    Xba I    NaeI

5'CATCTAGATGGCGCCGCCACAGTACGTTTGATCTCCAGCTTGGTCCC3' SEQ  
ID NO: 29 The V<sub>H</sub> primers, #2621 and #2853, extended from  
25 KpnI through PstI and were made to conserved regions of  
murine RNA.

#2621: 5'GGGGTACCAGGTCCA (A/G) CT (G/T) CTCGAGTC (A/T) GG3'  
SEQ ID NO: 30

30                    PstI  
#2853: 5'GCCTGCAGCTAGCTGAGGAGACGGTGACCGTGGTCCCTTGG-  
                         NheI

CCCCAG3' SEQ ID NO: 31

PCR, as described by Saiki et al., Science,  
35 239:487-491 (1988), was performed on the RNA template.  
For the PCR, the primers used were identified above. For

PCR amplification of  $V_H$ , DNA/primer mixtures consisted of 5  $\mu$ l RNA and 0.5  $\mu$ M of the primers. For PCR amplifications of  $V_L$ , DNA/primer mixtures consisted of 5  $\mu$ l RNA and 0.5  $\mu$ M of the primers. To these mixtures was added 250  $\mu$ M each of dATP, dCTP, dGTP and dTTP, 10 mM Tris-HCl pH 8.3, 50 mM KCl, 1.5 mM  $MgCl_2$ , 0.01% (w/v) gelatin, 0.01% (v/v) Tween 20, 0.01% (v/v) Nonidet P40 and 5 units AmpliTaq [Cetus]. Samples were subjected to 25 thermal cycles of PCR at 94°C, 30 seconds; 55°C, 30 seconds; 72°C, 45 seconds; ending with 5 minutes at 72°C. For cloning and sequencing, amplified  $V_H$  DNA was purified on a low melting point agarose gel and by Elutip-d column chromatography [Schleicher and Schuell-Dussel, Germany] and cloned into pUC18 [New England Biolabs]. The general cloning and ligation methodology was as described in Maniatis *et al.*, cited above.

$V_H$  DNA was cloned as KpnI-PstI fragments into similarly-digested pUC18.  $V_L$  DNA was cloned as HindIII-XbaI fragments into pUC18 digested with the same enzymes. Representative clones were sequenced by the dideoxy method [Sanger *et al.*, Proc. Natl. Acad. Sci. USA, 74:5463-5467 (1977)] using T7 DNA polymerase [US Biologicals]. From the sequences of NFS2  $V_H$  and  $V_L$  domains, the CDR sequences were elucidated in accordance with the methodology of Kabat *et al.*, in "Sequences of Proteins of Immunological Interest", US Dept of Health and Human Services, US Government Printing Office (1987) utilizing computer assisted alignment with other  $V_H$  and  $V_L$  sequences. The CDRs of the heavy and light chains of NFS2 are listed above and identified herein as SEQ ID NOS: 15-20 and 21-26.

Example 3 - Humanized Antibodies

The following example describes the preparation of an exemplary engineered antibody. Similar procedures may be followed for the development of engineered antibodies, using other *Plasmodium* antibodies or other anti-pathogen antibodies developed by conventional means.

The source of the donor CDRs utilized to prepare these engineered antibodies was the murine mAb, NFS2, described in Examples 1 and 2 above. The sequenced NFS2 variable framework regions were employed to again search through the Kabat database to identify homologous framework regions of a human antibody. The framework region of an antibody obtained from a human SLE patient B-cell hybridoma cell line 18/17 [H. Dersimonian et al., J. Immunol., 139:2496-2501 (1987)] was determined to be approximately 80% homologous to the NFS2 variable heavy chain framework region.

Given the murine NFS2 CDRs (Example 2) and the sequence of the human antibody 18/17, a synthetic heavy chain variable region was made, and PCR performed to fill in and amplify DNA. The NFS2 CDR sequences and the 18/17 V<sub>H</sub> framework regions were synthesized by the following overlapping oligonucleotides:

SEQ ID NO:44:TAGTGAAGAATTCGAGGACGCCAGCAACATGGTGTTCAGAC  
CCAGGTCTTCATTTCTCTGTTGCTCTGGATCTCTGGTGCCTACGGGGAGGTGCAG  
(Base 1-97);

SEQ ID NO:45:GCTAGCGGATTACCTTTAGCAGCCATGTCTCGGACCCCCCAGG  
GACTCTGAGAGGACACGTCGATCGCCTAAGTGGAATCCTATGCCATGAGCTGGG  
TCCGCCAGGCTCCAGGGAAAGGTCTAGAGTGGGTCTCAGAAATCTTTATAGTGAT  
GGTGGTAGTTAC (Base 158-259);

SEQ ID NO:46: GAACACGCTGTATCTGCAAATGAACAGCCTGAGAGCCGAG  
GACACGTCTCTGTTAAGGTTCTTGTGCGACATAGACGTTTACTGCAGTATATTAC  
TGTGCGAAACTCATCTACTATGGTTACGACGGGTATGCTATGGACTAGCTGCCCA  
TACGATACCTGATC (Base 316-421);

SEQ ID NO:47: TTCTTGAAAGCTTGGGCCCTTGGTACTAGCTGAGCTCACGG

TGACCAGGGTACCCTGGCCCCAGTAGTCCATAGCATACCCGTCG (Base 484-400);

SEQ ID NO:48: CATTTGCAGATACAGCGTGTTCCTTGAATTGTCTCTGGATA  
TCGTGAACCGGCCCGTCACAGTGTCTGGATAGTAGGTGTAACCTACCACCATCACT

5 AATTTC (Base 337-236);

SEQ ID NO:49: CTAAAGGTGAATCCGCTAGCTGCACAGGAGAGTCTCAGGGA  
CCCCCAGGCTGTACCAAGCCTCCCCCAGACTCGAGCAGCTGCACCTCCCCGTAG  
GCACC (Base 177-77).

10 These primers are annealed together and DNA is  
filled in using Taq polymerase, followed by PCR  
amplification using the following 5' primer:

SEQ ID NO:50: CCGCGAATTCGAGGACGCCAGCAAC

and 3' primer: SEQ ID NO:51:

CCGCAAGCTTGGGCCCTTGGTACTAGCT.

15 Any errors in the mapped sequence which were  
inserted by PCR were corrected. In addition,  
conservative nucleotide replacements were placed in the  
framework regions to introduce selected restriction sites  
suitable for enzymatic cleavage. These alterations in  
20 the framework regions are indicated by boxes in the  
sequences of Figs. 2 [SEQ ID NOS: 5 and 6], 3 [SEQ ID  
NOS: 7 and 8], 5 [SEQ ID NOS: 11 and 12] and 6 [SEQ ID  
NOS: 13 and 14]. Additionally, most murine and human  
antibodies have a basic residue before CDR3. Because the  
25 variable heavy chain of NFS2 has a non-basic residue Ser  
before CDR3 [SEQ ID NOS: 19-20 and 25 and 26], the  
acceptor basic residue (Lys) before CDR3 was deleted and  
replaced with Ser to create heavy chain Pfhzhc2-3.

30 Two synthetic heavy chain variable regions were  
obtained, namely, Pfhzhc2-3 SEQ ID NOS: 13 and 14, and  
Pfhzhc2-4 SEQ ID NOS: 11 and 12. These sequences are  
described in detail in Figs. 5 and 6. Each of these  
synthetic heavy chain variable regions is characterized  
by one or two nucleotide, or amino acid, differences.  
35 For example, Pfhzhc2-3 [SEQ ID NOS: 13 and 14] has a Ser



at position 98; and Pfhzhc2-4 [SEQ ID NOS: 11 and 12] has a Lys at position 98. Otherwise, these heavy chain variable regions are identical.

For a suitable light chain variable framework region, the NFS2 light chain CDRs and the light chain variable framework sequence of the human antibody identified in H. G. Klobeck *et al.*, Nucl. Acids Res., 13:6515-6529 (1985) were used to make a suitable synthetic light chain sequence by the same methods. The oligonucleotides used were as follows.

SEQ ID NO:52: TAAGCGGAATTCGTAGTCGGATATCGTGATGACCCAGTC  
TCCAGACTCGCTAGCTGTGTCTCTGGGCGAGAGGGC (Base 1-75);

SEQ ID NO:53: TTAAGTTGGCCTGGTATCAGCAGAAACCCGGGCGAGTCTCC  
TAAGTTGCTCATAGTTTTCTTAATGAACCGGACTTACTGGGCGTCAACTAG (Base  
130-198);

SEQ ID NO:54: GACAGATTTCACTCTCACCATCAGCAGCCTGCAGGCTGAA  
GATGTGGCAGTATACTACTGCTGTCTAAAGTGTCAGCAATATTATAGCTATCC  
(Base 241-321);

SEQ ID NO:55: CAGTTGGAAAGCTTGGCGCCGCCACAGTACGTTTGATCTCCA  
CCTTGGTCCCTCCGCCGAACGTCCGCGGATAGCTATAATATTGC (Base 389-  
304);

SEQ ID NO:56: GTGAAATCTGTCCCAGACCCGCTGCCACTGAATCGG  
TCAGGTACCCAGATTCCCTAGTTGACGCC (Base 252-187);

SEQ ID NO:57: CAGGCCAAGTAATTCTTTTGATTGCTCGAGTATAAA  
AGGCTCTGAGAGCTCTTGCAGTTGATGGTGGCCCTCTCGCCC (Base 141-64).

As described above, the primers were annealed together and DNA filled in using Taq polymerase, followed by PCR amplification with the following 5' SEQ ID NO:58: GCGGAATTCGTAGTCGGATATCGTGATGAC and 3' SEQ ID NO:59: TGGAAAGCTTGGCGCCGCCACAGTACGTTTGATC primers.

Two synthetic light chain variable sequences containing the NFS2 CDRs were designed and synthesized as described above for the synthetic heavy chains and referred to as Pfhzlc1-1 SEQ ID NOS: 5 and 6, and

Pfhzlc1-2 SEQ ID NOS: 7 and 8. These two sequences differed in amino acid sequence at only one amino acid position, 49. Pfhzlc1-1 [SEQ ID NOS: 5 and 6] has a Ser at position 49; Pfhzlc1-2 [SEQ ID NOS: 7 and 8] has a Pro at the same position.

These synthetic variable light and/or heavy chain sequences are employed in the construction of an exemplary humanized antibody. It is expected that any of the synthetic heavy chains will successfully associate with any of the synthetic light chains to produce a useful humanized antibody.

To produce a humanized antibody, for the heavy chain variable sequence Pfhzhc2-3 [SEQ ID NO: 13 and 14] (Fig. 6), the following signal sequence was synthesized onto this variable region: SEQ ID NOS: 60: ATGGTGTTCAG ACCCAGGTCTTCATTTCTCTGTTGCTCTGGATCTCTGGTGCCTAC, which encodes SEQ ID NO: 61: MetValLeuGlnThrGlnValPheIleSerLeuLeuLeuTrpIleSerGlyAlaTyr. For the synthetic light chain variable sequence Pfhzlc1-1 [SEQ ID NO: 5 and 6] (Fig. 2), the construct is digested with EcoRI and EcoRV, and the same signal sequence was ligated onto the variable sequence. Other signal sequences are well known to those of skill in the art and may be substituted for this exemplary sequence.

Selected constant regions of the human IgG<sub>1</sub> antibodies selected for the heavy and light chain were synthesized and confirmed by PCR. These constant region sequences were then inserted into pUC19-based expression vectors. The above-described synthetic variable constructs, containing the signal and variable regions of the light and heavy chains, were thereafter inserted into these pUC19-based expression vectors containing CMV promoters and the constant regions and fused in frame to the previously inserted human heavy and light chain

constant regions by conventional methods [Maniatis et al., cited above]. Thus, after insertion of the synthetic variable regions into these expression vectors, the plasmids shown in Figs. 7 and 8, resulted. These  
5 plasmids were then co-transfected into a selected host cell and, following incubation, the media was assayed for antibody activity via ELISA as described in Example 4 below.

Using similar techniques, another exemplary  
10 humanized antibody is constructed using the synthesized heavy chain sequence Pfhzhc2-3 [SEQ ID NO:13 and 14] (Fig. 6) and the synthetic light chain sequence Pfhzlc1-2 [SEQ ID NO:7 and 8].

#### Example 4 - A High Affinity Humanized Antibody

15 The amino acid differences in the variable regions of the frameworks of the original murine antibody NSF2 described in Examples 1 and 2 and the Pfhzhc2.3 were determined, and several changes were made to increase the level of conservation of the original antibody  
20 conformation.

At amino acid position 49, the Ser of the humanized heavy chain Pfhzhc2.3 was changed to Ala, which is the amino acid found at this position in the native murine NSF2. The replacement employed conventional  
25 genetic engineering technology, e.g., by making a synthetic DNA fragment containing the appropriate nucleotide changes to alter the amino acid. A fragment of Pfhzhc2.3 was digested with XbaI and EcoRV and the synthetic fragment bearing the nucleotide change encoding  
30 Ala in place of a Ser codon, is inserted to make the amino acid replacement. The resulting synthetic heavy chain was termed Pfhzhc2.6.

This synthetic heavy chain was expressed as previously described for the Pfhzhc2.3 synthetic heavy

chain. The expression plasmid for this humanized heavy chain sequence is essentially identical to the expression plasmid illustrated in Fig. 7, with the exception of the single amino acid difference described previously.

5           Similarly, humanized antibodies consisting of the Pfhzhc2.6 heavy chain and Pfhzlc1.1 light chain and the Pfhzhc2.6 heavy chain and the Pfhzlc1.2 light chain were assembled via co-transfection of mammalian cells and assayed for antibody activity by ELISA, as described in  
10   Example 5 below.

          Other high affinity antibodies specific for *P. falciparum* can be developed using a similar method designed to achieve minimal variable region framework modifications. The method involves the following order  
15   of steps of alteration and testing:

          (1) In addition to the alteration at amino acid position 49, other individual framework amino acid residues known to be critical for interaction with CDRs are compared in the primary antibody and the engineered  
20   CDR-replacement antibody. For example, heavy chain amino acid residue (Kabat numbering; see Kabat et al., cited above) is compared in the primary (donor) and engineered antibodies. A residue at this position is thought to interact with the invariant heavy chain CDR residue at  
25   position 94 (Lys-basic) via a salt bridge.

          If an amino acid is in the framework of the donor antibody but not in the framework of the engineered antibody, then an alternative heavy chain gene comprising the engineered antibody is produced. In the reverse  
30   situation whereby the engineered antibody framework comprises a residue at one position but the donor antibody does not, then an alternative heavy chain gene comprising the original amino acid at that position is reproduced. Prior to any further analysis, alternative

plasmids produced on this basis are tested for production of high affinity engineered antibodies.

(2) Framework amino acids within four residues of the CDRs as defined according to Kabat (see Kabat et al., cited above) are compared in the primary antibody and engineered CDR-replacement antibody. Where differences are present, then for each region the specific amino acids of that region are substituted for those in the corresponding region of the engineered antibody to provide a small number of engineered genes. Alternative plasmids produced on this basis are then tested for production of high affinity antibodies.

(3) Framework residues in the primary and engineered CDR-replacement antibodies are compared and residues with major differences in charge, size or hydrophobicity are highlighted. Alternative plasmids are produced on this basis with the individual highlighted amino acids represented by the corresponding amino acids of the primary antibody and such alternative plasmids are tested for production of high affinity antibodies.

#### Example 5: ELISA Assays

Expression of the synthetic heavy chain and light chain sequences were tested by transiently transfecting the plasmid DNAs into monkey COS cells. The following results are reported for Pfhzhc2-3 and Pfhzlc1-1. Ten micrograms of the plasmids are mixed together and ethanol precipitated. The DNAs are dissolved in Tris buffered saline (TBS) and mixed with DEAE-dextran (400 µg/ml final concentration)/chloroquine (0.1 mM), added to 3-4 x 10<sup>5</sup> COS cells grown in T25 flask, incubated for 4 hours at 37°C. Cells were shocked with 10% DMSO in phosphate buffered saline (PBS) for 1-2 minutes and after washing with PBS, cells were incubated in the presence of serum free growth medium. Media were collected 72 hours

post transfection (day 3 sample) and fresh media were added which were collected 120 hours post transfection (referred to as day 5 sample).

To compare the binding affinity of various antibodies, i.e., chimeric and humanized, large scale COS transfections were performed as described above. For each antibody, 200  $\mu$ g heavy chain plasmid and 200  $\mu$ g light chain plasmid DNAs were used to transfect a total of  $2.5 \times 10^7$  COS cells. The media collected (day 3 and day 5) were pooled, assayed for antibody expression using  $F_c$  capture ELISA. The media were concentrated using Amicon to 6 ml. Amount of antibody in the pooled media varied from 9 mg/ml to 25 mg/ml. These concentrated samples were used to compare binding affinities via the ISI and ILSDA assays.

The presence of humanized antibody in the medium of wells containing transfected clones is measured by conventional ELISA techniques. Micro-titer plates are coated overnight at 4°C with goat anti-human IgG ( $F_c$  specific) antibodies [Sigma, St. Louis, MO] at 0.1  $\mu$ g per well. After washing with PBS (pH 7.5), 50  $\mu$ l of culture medium from the wells containing transfectants is added to each microtitre well for 2 hours at room temperature. The wells are then emptied, washed with PBS and peroxidase-conjugated goat anti-human IgG antibodies [BioRad, Richmond, CA] are added at 50  $\mu$ L of a 1/1000 dilution per well. Plates are then incubated at room temperature for 1 hour. The wells are then emptied and washed with PBS. 100  $\mu$ l 2,2'-azino-di[3-ethyl-benzthiazoline sulfonate(6)] are added per well. Reactions were allowed to continue for 1 hour at room temperature. The absorbance at 405 nm is then measured spectrophotometrically. The ability of the humanized antibody in the medium of wells containing transfected clones to bind *P. falciparum* circumsporozoite protein was

measured by ELISA. Microtiter plates were coated overnight at 4°C with *E. coli*-produced R32tet32 at 0.1 µg per well. After washing with PBS, 50 µl of culture medium from the wells which contain transfectants is added to each microtiter well for 2 hours at room temperature. The wells are then emptied, washed with PBS and peroxidase-conjugated goat anti-human IgG antibodies [BioRad, Richmond, CA] are added at 50 µL of a 1/1000 dilution per well. Plates are then incubated at room temperature for 1 hour. The wells are then emptied and washed with PBS. 100 µl 2,2'-azino-di[3-ethyl-benzthiazoline sulfonate(6)] are added per well. Reactions were allowed to continue for 1 hour at room temperature. The absorbance at 405 nm is then measured spectrophotometrically.

In preliminary studies, an increase in affinity was observed for the humanized antibodies of Example 4, which contained the Pfhzhc2-6 heavy chain construct, as compared to the Pfhzhc2-3 heavy chain construct.

#### 20 Example 6 - Construction of Chimeric Antibody

A chimeric antibody of the invention was constructed essentially as described above. A chimeric antibody contains the native murine NSF2 variable framework and CDR regions on the human constant regions selected for the heavy chain [H. Dersimonian et al., J. Immunol., 139:2496-2501 (1987) and light chain [Klobeck et al., Nucl. Acids. Res., 13:6515-6529 (1985)], with the exception that the variable regions were obtained by PCR of the RNA of the murine antibody obtained from the NFS2 hybridoma and the entire constant regions of the human IgG<sub>1</sub> antibodies were synthesized by overlapping oligonucleotides and amplified by PCR. Any errors which were inserted by PCR were corrected. The resulting

chimeric heavy chain and chimeric light chain were expressed as described above for the humanized antibody.

This chimeric antibody is advantageous in that it is characterized by activity substantially identical to that of the native murine antibody, but contains  
5 enough human sequences that it is anticipated to be useful in human therapy.

#### Example 7 - ISI Assay

The Inhibition of Sporozoite Invasion assay is  
10 performed as described in M. R. Hollingdale et al., J. Immunol., 132:909-913 (1984) to be used to assess neutralizing effect against live *P. falciparum* sporozoites. In the ISI assay, the human hepatoma cloned cell line HepG2-A16<sub>2</sub> was grown to near confluency on 1%  
15 CO<sub>2</sub> glass cover slips in MEM and 10% bovine fetal serum. Antisera or purified antibodies were diluted in culture medium (see table below) and added to the cell cultures. 30,000 *P. falciparum* sporozoites isolated from dissected mosquito salivary glands are counted, diluted and added  
20 to each cell culture. The cultures are incubated at 37°C for 2.5 hours, rinsed with PES, and fixed with methanol. Fixed cultures are reacted in an immunoperoxidase antibody assay using a labelled mAb which recognizes the *P. falciparum* CS protein to visualize invaded  
25 sporozoites. Then, the number of invading sporozoites are counted by phase microscopy at 400x. The ISI is the percent reduction in invasion in the presence of the test antibody, the humanized antibody, as compared to a control (i.e. non-related) antibody. The assay ranks  
30 antibodies in order according to their relative potency.

The following table provides the results performed on the chimeric and synthetic antibodies described previously. Values given are percent inhibition and are the average of 2-3 independent assays.



---

Summary of ISI Studies

	In $\mu\text{g/ml}$ :	<u>20</u>	<u>10</u>	<u>5.0</u>	<u>2.0</u>	<u>1.0</u>	<u>0.1</u>
	Chimeric	99	98.5	98	88	83	50
5	PfHzhc2-3/lc1-1	92	75.5		60		
	PfHzhc2-3/lc1-2	85	80.0		0		
	PfHzhc2-6/lc1-1		90.0	75		53	0
	PfHzhc2-6/lc1-2		87.0	65		50	0

---

- 10                    Numerous modifications and variations of the present invention are included in the above-identified specification and are expected to be obvious to one of skill in the art. For example, recombinant antibodies capable of neutralizing pathogens other than *P.*
- 15    *falciparum* may be provided according to the teachings of this invention for the development of prophylactic agents capable of conferring passive immunity to other human diseases. Preferably, engineered antibodies capable of recognizing repeat regions on other malaria pathogens or
- 20    engineered antibodies to any region on the surface of any stages of the life-cycle of the plasmodium species or capable of neutralizing any stage in the life cycle of the parasite, are desirable starting materials to develop passive immunity agents according to this invention.
- 25    Such modifications and alterations to the compositions and processes of the present invention are believed to be encompassed in the scope of the claims appended hereto.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: SmithKline Beecham, Corporation  
U. S. Government, Secretary of  
the Navy  
U. S. Government, Secretary of  
the Army
- (ii) TITLE OF INVENTION: Novel Antibodies for Conferring  
Passive Immunity Against Infection by a  
Pathogen in Man
- (iii) NUMBER OF SEQUENCES: 61
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: Howson and Howson
  - (B) STREET: Box 457, 321 Norristown Road
  - (C) CITY: Spring House
  - (D) STATE: PA
  - (E) COUNTRY: USA
  - (F) ZIP: 19477
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk
  - (B) COMPUTER: IBM PC compatible
  - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
  - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER:
  - (B) FILING DATE:
  - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: US 07/941,654
  - (B) FILING DATE: 09-SEP-1992
- (viii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: Bak, Mary E.
  - (B) REGISTRATION NUMBER: 31,215
  - (C) REFERENCE/DOCKET NUMBER: SBC P50107
- (ix) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: (215) 540-9200
  - (B) TELEFAX: (215) 540-5818

49

## (2) INFORMATION FOR SEQ ID NO:1:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 164 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: unknown

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

```

Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn
 1              5              10              15
Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala
          20              25              30
Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn
          35              40              45
Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro
          50              55              60
Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn
          65              70              75
Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala
          80              85              90
Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn
          95              100             105
Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro
          110             115             120
Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn
          125             130             135
Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Ala Asn Pro Asn Val
          140             145             150
Asp Pro Asn Val Asp Pro Asn Val Asp Pro Asn Val Asp Pro
          155             160

```

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 163 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: unknown

50

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met	Asp	Pro	Asn	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	Ala	Asn	Pro	1	5	10	15
Asn	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	Ala	Asn	20	25	30	
Pro	Asn	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	Ala	35	40	45	
Asn	Pro	Asn	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	50	55	60	
Ala	Asn	Pro	Asn	Val	Asp	Pro	Asn	Ala	Asn	Pro	Asn	Ala	Asn	Pro	65	70	75	
Asn	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	Ala	Asn	80	85	90	
Pro	Asn	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	Ala	95	100	105	
Asn	Pro	Asn	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	110	115	120	
Ala	Asn	Pro	Asn	Ala	Asn	Pro	Asn	Val	Asp	Pro	Leu	Arg	Arg	Thr	125	130	135	
His	Arg	Gly	Arg	His	His	Arg	Arg	His	Arg	Cys	Gly	Cys	Trp	Arg	140	145	150	
Leu	Tyr	Arg	Arg	His	His	Arg	Trp	Gly	Arg	Ser	Gly	Ser	155	160				

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 339 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..339

51

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GAT	ATT	CAG	CTG	ACC	CAG	TCT	CCA	TCC	TCC	CTA	GCT	GTG	TCA	42
Asp	Ile	Gln	Leu	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ala	Val	Ser	
1				5					10					
GTT	GGA	GAG	AAG	GTT	ACT	ATG	AGC	TGC	AAG	TCC	AGT	CAG	AGC	84
Val	Gly	Glu	Lys	Val	Thr	Met	Ser	Cys	Lys	Ser	Ser	Gln	Ser	
15					20					25				
CTT	TTA	TAT	AGT	AGC	AAT	CAA	AAG	AAT	TAC	TTG	GCC	TGG	TAC	126
Leu	Leu	Tyr	Ser	Ser	Asn	Gln	Lys	Asn	Tyr	Leu	Ala	Trp	Tyr	
	30					35					40			
CAG	CAG	AAA	CCA	GGG	CAG	TCT	CCT	AAA	CTG	CTG	ATT	TAC	TGG	168
Gln	Gln	Lys	Pro	Gly	Gln	Ser	Pro	Lys	Leu	Leu	Ile	Tyr	Trp	
		45					50					55		
GCA	TCC	ACT	AGG	GAA	TCT	GGG	GTC	CCT	GAT	CGC	TTC	ACA	GGC	210
Ala	Ser	Thr	Arg	Glu	Ser	Gly	Val	Pro	Asp	Arg	Phe	Thr	Gly	
			60					65					70	
AGA	GGA	TCC	GGG	ACA	GAT	TTC	ACT	CTC	ACC	ATC	AGC	AGT	GTG	252
Arg	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Val	
				75					80					
AAG	GCT	GAA	GAC	CTG	GCA	GTT	TAT	TAC	TGT	CAG	CAA	TAT	TAT	294
Lys	Ala	Glu	Asp	Leu	Ala	Val	Tyr	Tyr	Cys	Gln	Gln	Tyr	Tyr	
85					90					95				
AGC	TAT	CCT	CGG	ACG	TTC	GGT	GGA	GGG	ACC	AAG	CTG	GAG	ATC	336
Ser	Tyr	Pro	Arg	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	
	100					105					110			
AAA														339
Lys														

## (2) INFORMATION FOR SEQ ID NO:4:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 113 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Asp	Ile	Gln	Leu	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ala	Val	Ser	Val
1				5					10					15

52

Gly	Glu	Lys	Val	Thr	Met	Ser	Cys	Lys	Ser	Ser	Gln	Ser	Leu	Leu	
				20					25					30	
Tyr	Ser	Ser	Asn	Gln	Lys	Asn	Tyr	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	
				35					40					45	
Pro	Gly	Gln	Ser	Pro	Lys	Leu	Leu	Ile	Tyr	Trp	Ala	Ser	Thr	Arg	
				50					55					60	
Glu	Ser	Gly	Val	Pro	Asp	Arg	Phe	Thr	Gly	Arg	Gly	Ser	Gly	Thr	
				65					70					75	
Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Val	Lys	Ala	Glu	Asp	Leu	Ala	
				80					85					90	
Val	Tyr	Tyr	Cys	Gln	Gln	Tyr	Tyr	Ser	Tyr	Pro	Arg	Thr	Phe	Gly	
				95					100					105	
Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys								
				110											

## (2) INFORMATION FOR SEQ ID NO:5:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 339 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

## (ii) MOLECULE TYPE: DNA (genomic)

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..339

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

GAT	ATC	GTG	ATG	ACC	CAG	TCT	CCA	GAC	TCG	CTA	GCT	GTG	TCT	42
Asp	Ile	Val	Met	Thr	Gln	Ser	Pro	Asp	Ser	Leu	Ala	Val	Ser	
1				5					10					
CTG	GGC	GAG	AGG	GCC	ACC	ATC	AAC	TGC	AAG	AGC	TCT	CAG	AGC	84
Leu	Gly	Glu	Arg	Ala	Thr	Ile	Asn	Cys	Lys	Ser	Ser	Gln	Ser	
15				20					25					
CTT	TTA	TAC	TCG	AGC	AAT	CAA	AAG	AAT	TAC	TTG	GCC	TGG	TAT	126
Leu	Leu	Tyr	Ser	Ser	Asn	Gln	Lys	Asn	Tyr	Leu	Ala	Trp	Tyr	
30					35						40			

53

CAG	CAG	AAA	CCC	GGG	CAG	TCT	CCT	AAG	TTG	CTC	ATT	TAC	TGG	168
Gln	Gln	Lys	Pro	Gly	Gln	Ser	Pro	Lys	Leu	Leu	Ile	Tyr	Trp	
		45					50					55		
GCG	TCA	ACT	AGG	GAA	TCT	GGG	GTA	CCT	GAC	CGA	TTC	AGT	GGC	210
Ala	Ser	Thr	Arg	Glu	Ser	Gly	Val	Pro	Asp	Arg	Phe	Ser	Gly	
			60					65					70	
AGC	GGG	TCT	GGG	ACA	GAT	TTC	ACT	CTC	ACC	ATC	AGC	AGC	CTG	252
Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	
				75				80						
CAG	GCT	GAA	GAT	GTG	GCA	GTA	TAC	TAC	TGT	CAG	CAA	TAT	TAT	294
Gln	Ala	Glu	Asp	Val	Ala	Val	Tyr	Tyr	Cys	Gln	Gln	Tyr	Tyr	
85					90					95				
AGC	TAT	CCG	CGG	ACG	TTC	GGC	GGA	GGG	ACC	AAG	GTG	GAG	ATC	336
Ser	Tyr	Pro	Arg	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Val	Glu	Ile	
	100					105					110			
AAA														339
Lys														

## (2) INFORMATION FOR SEQ ID NO:6:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 113 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Asp	Ile	Val	Met	Thr	Gln	Ser	Pro	Asp	Ser	Leu	Ala	Val	Ser	Leu
1				5					10					15
Gly	Glu	Arg	Ala	Thr	Ile	Asn	Cys	Lys	Ser	Ser	Gln	Ser	Leu	Leu
				20					25					30
Tyr	Ser	Ser	Asn	Gln	Lys	Asn	Tyr	Leu	Ala	Trp	Tyr	Gln	Gln	Lys
				35					40					45
Pro	Gly	Gln	Ser	Pro	Lys	Leu	Leu	Ile	Tyr	Trp	Ala	Ser	Thr	Arg
				50					55					60
Glu	Ser	Gly	Val	Pro	Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr
				65					70					75

54

```

Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Ala Glu Asp Val Ala
      80                      85                      90
Val Tyr Tyr Cys Gln Gln Tyr Tyr Ser Tyr Pro Arg Thr Phe Gly
      95                      100                     105
Gly Gly Thr Lys Val Glu Ile Lys
      110

```

## (2) INFORMATION FOR SEQ ID NO:7:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 339 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

## (ii) MOLECULE TYPE: DNA (genomic)

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..339

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

```

GAT ATC GTG ATG ACC CAG TCT CCA GAC TCG CTA GCT GTG TCT      42
Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser
  1                      5                      10

CTG GGC GAG AGG GCC ACC ATC AAC TGC AAG AGC TCT CAG AGC      84
Leu Gly Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser
 15                      20                      25

CTT TTA TAC TCG AGC AAT CAA AAG AAT TAC TTG GCC TGG TAT     126
Leu Leu Tyr Ser Ser Asn Gln Lys Asn Tyr Leu Ala Trp Tyr
 30                      35                      40

CAG CAG AAA CCC GGG CAG CCT CCT AAG TTG CTC ATT TAC TGG     168
Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile Tyr Trp
 45                      50                      55

GCG TCG ACT AGG GAA TCT GGG GTA CCT GAC CGA TTC AGT GGC     210
Ala Ser Thr Arg Glu Ser Gly Val Pro Asp Arg Phe Ser Gly
 60                      65                      70

AGC GGG TCT GGG ACA GAT TTC ACT CTC ACC ATC AGC AGC CTG     252
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu
 75                      80

```



[illegible]

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 113 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Asp 1	Ile	Val	Met	Thr 5	Gln	Ser	Pro	Asp	Ser 10	Leu	Ala	Val	Ser	Leu 15
Gly	Glu	Arg	Ala	Thr 20	Ile	Asn	Cys	Lys	Ser 25	Ser	Gln	Ser	Leu	Leu 30
Tyr	Ser	Ser	Asn	Gln 35	Lys	Asn	Tyr	Leu	Ala 40	Trp	Tyr	Gln	Gln	Lys 45
Pro	Gly	Gln	Pro	Pro 50	Lys	Leu	Leu	Ile	Tyr 55	Trp	Ala	Ser	Thr	Arg 60
Glu	Ser	Gly	Val	Pro 65	Asp	Arg	Phe	Ser	Gly 70	Ser	Gly	Ser	Gly	Thr 75
Asp	Phe	Thr	Leu	Thr 80	Ile	Ser	Ser	Leu	Gln 85	Ala	Glu	Asp	Val	Ala 90
Val	Tyr	Tyr	Cys	Gln 95	Gln	Tyr	Tyr	Ser	Tyr 100	Pro	Arg	Thr	Phe	Gly 105
Gly	Gly	Thr	Lys	Val 110	Glu	Ile	Lys							



## (2) INFORMATION FOR SEQ ID NO:10:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 118 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Leu	Glu	Ser	Gly	Gly	Gly	Leu	Val	Lys	Pro	Gly	Gly	Ser	Leu	Lys	1	5	10	15
Ile	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Ser	Tyr	Ala	Met	20	25	30	
Ser	Trp	Val	Arg	Gln	Ser	Pro	Glu	Lys	Arg	Leu	Glu	Trp	Val	Ala	35	40	45	
Glu	Ile	Ser	Asp	Gly	Gly	Ser	Tyr	Thr	Tyr	Tyr	Pro	Asp	Thr	Val	50	55	60	
Thr	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ala	Lys	Asn	Thr	Leu	65	70	75	
Tyr	Leu	Glu	Met	Ser	Ser	Leu	Arg	Ser	Glu	Asp	Thr	Ala	Met	Tyr	80	85	90	
Tyr	Cys	Ala	Ser	Leu	Ile	Tyr	Tyr	Gly	Tyr	Asp	Gly	Tyr	Ala	Met	95	100	105	
Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Ser	Val	Thr	Val	Ser	Ser	110	115				

## (2) INFORMATION FOR SEQ ID NO:11:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 389 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

## (ii) MOLECULE TYPE: DNA (genomic)

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..366

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

GAG	GTG	CAG	CTG	CTC	GAG	TCT	GGG	GGA	GGC	TTG	GTA	CAG	CCT	42
Glu	Val	Gln	Leu	Leu	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	
1				5					10					
GGG	GGG	TCC	CTG	AGA	CTC	TCC	TGT	GCA	GCT	AGC	GGA	TTC	ACC	84
Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	
15				20						25				
TTT	AGC	AGC	TAT	GCC	ATG	AGC	TGG	GTC	CGC	CAG	GCT	CCA	GGG	126
Phe	Ser	Ser	Tyr	Ala	Met	Ser	Trp	Val	Arg	Gln	Ala	Pro	Gly	
30						35					40			
AAA	GGT	CTA	GAG	TGG	GTC	TCA	GAA	ATT	AGT	GAT	GGT	GGT	AGT	168
Lys	Gly	Leu	Glu	Trp	Val	Ser	Glu	Ile	Ser	Asp	Gly	Gly	Ser	
		45					50					55		
TAC	ACC	TAC	TAT	CCA	GAC	ACT	GTG	ACG	GGC	CGG	TTC	ACG	ATA	210
Tyr	Thr	Tyr	Tyr	Pro	Asp	Thr	Val	Thr	Gly	Arg	Phe	Thr	Ile	
			60					65					70	
TCC	AGA	GAC	AAT	TCC	AAG	AAC	ACG	CTG	TAT	CTG	CAA	ATG	AAC	252
Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr	Leu	Gln	Met	Asn	
				75					80					
AGC	CTG	AGA	GCC	GAG	GAC	ACT	GCA	GTA	TAT	TAC	TGT	GCG	AAA	294
Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Ala	Lys	
85					90					95				
CTC	ATC	TAC	TAT	GGT	TAC	GAC	GGG	TAT	GCT	ATG	GAC	TAC	TGG	336
Leu	Ile	Tyr	Tyr	Gly	Tyr	Asp	Gly	Tyr	Ala	Met	Asp	Tyr	Trp	
100						105						110		
GGC	CAG	GGT	ACC	CTG	GTC	ACC	GTG	AGC	TCA	GCTAGTACCA				376
Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser					
		115					120							
AGGGCCCAAG	CTT													389

## (2) INFORMATION FOR SEQ ID NO:12:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 122 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

59

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Glu	Val	Gln	Leu	Leu	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	
1				5					10					15	
Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	Ser	
				20					25					30	
Ser	Tyr	Ala	Met	Ser	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	
				35					40					45	
Glu	Trp	Val	Ser	Glu	Ile	Ser	Asp	Gly	Gly	Ser	Tyr	Thr	Tyr	Tyr	
				50					55					60	
Pro	Asp	Thr	Val	Thr	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	
				65					70					75	
Lys	Asn	Thr	Leu	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	
				80					85					90	
Thr	Ala	Val	Tyr	Tyr	Cys	Ala	Lys	Leu	Ile	Tyr	Tyr	Gly	Tyr	Asp	
				95					100					105	
Gly	Tyr	Ala	Met	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	
				110					115					120	
Ser	Ser														

## (2) INFORMATION FOR SEQ ID NO:13:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 389 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

## (ii) MOLECULE TYPE: DNA (genomic)

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..366

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GAG	GTG	CAG	CTG	CTC	GAG	TCT	GGG	GGA	GGC	TTG	GTA	CAG	CCT		42
Glu	Val	Gln	Leu	Leu	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro		
1				5					10						
GGG	GGG	TCC	CTG	AGA	CTC	TCC	TGT	GCA	GCT	AGC	GGA	TTC	ACC		84
Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr		
15				20					25						

60

TTT	AGC	AGC	TAT	GCC	ATG	AGC	TGG	GTC	CGC	CAG	GCT	CCA	GGG	126
Phe	Ser	Ser	Tyr	Ala	Met	Ser	Trp	Val	Arg	Gln	Ala	Pro	Gly	
	30					35					40			
AAA	GGT	CTA	GAG	TGG	GTC	TCA	GAA	ATT	AGT	GAT	GGT	GGT	AGT	168
Lys	Gly	Leu	Glu	Trp	Val	Ser	Glu	Ile	Ser	Asp	Gly	Gly	Ser	
		45					50					55		
TAC	ACC	TAC	TAT	CCA	GAC	ACT	GTG	ACG	GGC	CGG	TTC	ACG	ATA	210
Tyr	Thr	Tyr	Tyr	Pro	Asp	Thr	Val	Thr	Gly	Arg	Phe	Thr	Ile	
			60					65					70	
TCC	AGA	GAC	AAT	TCC	AAG	AAC	ACG	CTG	TAT	CTG	CAA	ATG	AAC	252
Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr	Leu	Gln	Met	Asn	
				75					80					
AGC	CTG	AGA	GCC	GAG	GAC	ACT	GCA	GTG	TAT	TAC	TGT	GCA	TCT	294
Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Ala	Ser	
	85				90					95				
CTC	ATC	TAC	TAT	GGT	TAC	GAC	GGG	TAT	GCT	ATG	GAC	TAC	TGG	336
Leu	Ile	Tyr	Tyr	Gly	Tyr	Asp	Gly	Tyr	Ala	Met	Asp	Tyr	Trp	
	100					105					110			
GGC	CAA	GGT	ACC	CTG	GTC	ACC	GTG	AGC	TCA	GCTAGTACCA				376
Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser					
		115					120							
AGGGCCCAAG CTT														389

## (2) INFORMATION FOR SEQ ID NO:14:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 122 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Glu	Val	Gln	Leu	Leu	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly
1				5					10					15
Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	Ser
				20					25					30
Ser	Tyr	Ala	Met	Ser	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu
				35					40					45

61

Glu	Trp	Val	Ser	Glu	Ile	Ser	Asp	Gly	Gly	Ser	Tyr	Thr	Tyr	Tyr
				50					55					60
Pro	Asp	Thr	Val	Thr	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser
				65					70					75
Lys	Asn	Thr	Leu	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp
				80					85					90
Thr	Ala	Val	Tyr	Tyr	Cys	Ala	Ser	Leu	Ile	Tyr	Tyr	Gly	Tyr	Asp
				95					100					105
Gly	Tyr	Ala	Met	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val
				110					115					120
Ser	Ser													

## (2) INFORMATION FOR SEQ ID NO:15:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

AGCTATGCCA TGAGC

15

## (2) INFORMATION FOR SEQ ID NO:16:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: unknown

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Ser	Tyr	Ala	Met	Ser
1				5

62

## (2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 51 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

GAAATTAGTG ATGGTGGTAG TTACACCTAC TATCCAGACA CTGTGACGGG C 51

## (2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 17 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Glu Ile Ser Asp Gly Gly Ser Tyr Thr Tyr Tyr Pro Asp Thr  
1 5 10  
Val Thr Gly  
15

## (2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 39 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

CTCATCTACT ATGGTTACGA CGGGTATGCT ATGGACTAC 39



- (ii) MOLECULE TYPE: protein

Leu Ile Tyr Tyr Gly Tyr Asp Gly Tyr Ala Met Asp Tyr  
1 5 10

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 51 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: unknown

AAGAGCTCTC AGAGCCTTTT ATACTCGAGC AATCAAAGA ATTACTTGGC C 51

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 17 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: unknown

Lys Ser Ser Gln Ser Leu Leu Tyr Ser Ser Asn Gln Lys Asn  
1 5 10

Tyr Leu Ala  
15

64

## (2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 21 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

TGGGCGTCAA CTAGGGAATC T

21

## (2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 7 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

Trp	Ala	Ser	Thr	Arg	Glu	Ser
1				5		

## (2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 27 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

CAGCAATATT ATAGCTATCC GCGGACG

27

## (2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 9 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: unknown

65

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

Gln Gln Tyr Tyr Ser Tyr Pro Arg Thr  
1 5

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 6 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

Pro Asn Ala Asn Pro Asn  
1 5

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

CCAGATGTAA GCTTCAGCTG ACCCAGTCTC CA

32

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 47 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

66

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

CATCTAGATG GCGCCGCCAC AGTACGTTTG ATCTCCAGCT TGGTCCC

47

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

GGGGTACCAG GTCCARCTKC TCGAGTCWGG

30

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 47 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

GCCTGCAGCT AGCTGAGGAG ACGGTGACCG TGGTCCCTTG GCCCCAG

47

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

AGCTATGCCA TGTCT

15

67

## (2) INFORMATION FOR SEQ ID NO:33:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 5 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

Ser Tyr Ala Met Ser  
1 5

## (2) INFORMATION FOR SEQ ID NO:34:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 51 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

AAGTCCAGTC AGAGCCTTTT ATATAGTAGC AATCAAAGA ATTACTTGGC C 51

## (2) INFORMATION FOR SEQ ID NO:35:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 17 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

Lys Ser Ser Gln Ser Leu Leu Tyr Ser Ser Asn Gln Lys Asn  
1 5 10

Tyr Leu Ala  
15

68

## (2) INFORMATION FOR SEQ ID NO:36:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 21 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

TGGGCATCCA CTAGGGAATC T

21

## (2) INFORMATION FOR SEQ ID NO:37:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 7 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

Trp	Ala	Ser	Thr	Arg	Glu	Ser
1				5		

## (2) INFORMATION FOR SEQ ID NO:38:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 27 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

CAGCAATATT ATAGCTATCC TCGGACG

27

69

## (2) INFORMATION FOR SEQ ID NO:39:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 9 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

Gln Gln Tyr Tyr Ser Tyr Pro Arg Thr  
1 5

## (2) INFORMATION FOR SEQ ID NO:40:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 21 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

TGGGCGTCGA CTAGGGAATC T

21

## (2) INFORMATION FOR SEQ ID NO:41:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 7 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

Trp Ala Ser Thr Arg Glu Ser  
1 5

70

## (2) INFORMATION FOR SEQ ID NO:42:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 366 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: unknown

## (ii) MOLECULE TYPE: DNA (genomic)

## (ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 1..366

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

GAG	GTG	CAG	CTG	CTC	GAG	TCT	GGG	GGA	GGC	TTG	GTA	CAG	CCT	42
Glu	Val	Gln	Leu	Leu	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	
1				5					10					
GGG	GGG	TCC	CTG	AGA	CTC	TCC	TGT	GCA	GCT	AGC	GGA	TTC	ACC	84
Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	
15				20					25					
TTT	AGC	AGC	TAT	GCC	ATG	AGC	TGG	GTC	CGC	CAG	GCT	CCA	GGG	126
Phe	Ser	Ser	Tyr	Ala	Met	Ser	Trp	Val	Arg	Gln	Ala	Pro	Gly	
30				35					40					
AAA	GGT	CTA	GAG	TGG	GTC	GCA	GAG	ATC	TCT	GAT	GGT	GGT	AGT	168
Lys	Gly	Leu	Glu	Trp	Val	Ala	Glu	Ile	Ser	Asp	Gly	Gly	Ser	
		45					50				55			
TAC	ACC	TAC	TAT	CCA	GAC	ACT	GTG	ACG	GGC	CGG	TTC	ACG	ATA	210
Tyr	Thr	Tyr	Tyr	Pro	Asp	Thr	Val	Thr	Gly	Arg	Phe	Thr	Ile	
			60				65						70	
TCC	AGA	GAC	AAT	TCC	AAG	AAC	ACG	CTG	TAT	CTG	CAA	ATG	AAC	252
Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr	Leu	Gln	Met	Asn	
			75						80					
AGC	CTG	AGA	GCC	GAG	GAC	ACT	GCA	GTG	TAT	TAC	TGT	GCA	TCT	294
Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Ala	Ser	
85				90					95					
CTC	ATC	TAC	TAT	GGT	TAC	GAC	GGG	TAT	GCT	ATG	GAC	TAC	TGG	336
Leu	Ile	Tyr	Tyr	Gly	Tyr	Asp	Gly	Tyr	Ala	Met	Asp	Tyr	Trp	
100				105							110			
GGC	CAA	GGT	ACC	CTG	GTC	ACC	GTG	AGC	TCA					366
Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser					
		115					120							



71

## (2) INFORMATION FOR SEQ ID NO:43:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 122 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

Glu	Val	Gln	Leu	Leu	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	1	5	10	15
Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	Ser	20	25	30	
Ser	Tyr	Ala	Met	Ser	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	35	40	45	
Glu	Trp	Val	Ala	Glu	Ile	Ser	Asp	Gly	Gly	Ser	Tyr	Thr	Tyr	Tyr	50	55	60	
Pro	Asp	Thr	Val	Thr	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	65	70	75	
Lys	Asn	Thr	Leu	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	80	85	90	
Thr	Ala	Val	Tyr	Tyr	Cys	Ala	Ser	Leu	Ile	Tyr	Tyr	Gly	Tyr	Asp	95	100	105	
Gly	Tyr	Ala	Met	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	110	115	120	
Ser	Ser																	

## (2) INFORMATION FOR SEQ ID NO:44:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 97 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

TAGTGAAGAA TTCGAGGACG CCAGCAACAT GGTGTTGCAG ACCCAGGTCT	50
TCATTTCTCT GTTGCTCTGG ATCTCTGGTG CCTACGGGGA GGTGCAG	97

## (2) INFORMATION FOR SEQ ID NO:45:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 164 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

GCTAGCGGAT TCACCTTTAG CAGCCATGTC GGACCCCCCA GGGACTCTGA	50
GAGGACACGT CGATCGCCTA AGTGGAATC CTATGCCATG AGCTGGGTCC	100
GCCAGGCTCC AGGGAAAGGT CTAGAGTGGG TCTCAGAAAT CTTTATAGTG	150
ATGGTGGTAG TTAC	164

## (2) INFORMATION FOR SEQ ID NO:46:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 164 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

GAACACGCTG TATCTGCAAA TGAACAGCCT GAGAGCCGAG GACACGTCTC	50
TGTTAAGGTT CTTGTGCGAC ATAGACGTTT ACTGCAGTAT ATTACTGTGC	100
GAAACTCATC TACTATGGTT ACGACGGGTA TGCTATGGAC TAGCTGCCCA	150
TACGATACCT GATC	164

## (2) INFORMATION FOR SEQ ID NO:47:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 85 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

TTCTTGAAAG CTTGGGCCCT TGGTACTAGC TGAGCTCACC GTGACCAGGG	50
TACCCTGGCC CCAGTAGTCC ATAGCATACC CGTCG	85

## (2) INFORMATION FOR SEQ ID NO:48:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 102 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

CATTTGCAGA TACAGCGTGT TCTTGGAATT GTCTCTGGAT ATCGTGAACC	50
GGCCCGTCAC AGTGTCTGGA TAGTAGGTGT AACTACCACC ATCACTAATT TC	102

## (2) INFORMATION FOR SEQ ID NO:49:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 101 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

CTAAAGGTGA ATCCGCTAGC TGCACAGGAG AGTCTCAGGG ACCCCCCAGG	50
CTGTACCAAG CCTCCCCCAG ACTCGAGCAG CTGCACCTCC CCGTAGGCAC C	101

74

## (2) INFORMATION FOR SEQ ID NO:50:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 25 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

CCGCGAATTC GAGGACGCCA GCAAC

25

## (2) INFORMATION FOR SEQ ID NO:51:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 28 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

CCGCAAGCTT GGGCCCTTGG TACTAGCT

28

## (2) INFORMATION FOR SEQ ID NO:52:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 75 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

TAAGCGGAAT TCGTAGTCGG ATATCGTGAT GACCCAGTCT CCAGACTCGC

50

TAGCTGTGTC TCTGGGCGAG AGGGC

75

## (2) INFORMATION FOR SEQ ID NO:53:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 90 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

TTACTTGGCC TGGTATCAGC AGAAACCCGG GCAGTCTCCT AAGTTGCTCA	50
TAGTTTTCTT AATGAACCGG ACTTACTGGG CGTCAACTAG	90

## (2) INFORMATION FOR SEQ ID NO:54:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 93 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

GACAGATTTT ACTCTCACCA TCAGCAGCCT GCAGGCTGAA GATGTGGCAG	50
TATACTACTG CTGTCTAAAG TGTCAGCAAT ATTATAGCTA TCC	93

## (2) INFORMATION FOR SEQ ID NO:55:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 86 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

CAGTTGGAAG GCTTGGCGCC GCCACAGTAC GTTTGATCTC CACCTTGGTC 50  
CCTCCGCCGA ACGTCCGCGG ATAGCTATAA TATTGC 86

## (2) INFORMATION FOR SEQ ID NO:56:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 66 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: unknown

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

GTGAAATCTG TCCCAGACCC GCTGCCACTG AATCGGTCAG GTACCCCAGA 50  
TTCCCTAGTT GACGCC 66

## (2) INFORMATION FOR SEQ ID NO:57:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 78 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: unknown

## (ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

CAGGCCAAGT AATTCTTTTG ATTGCTCGAG TATAAAAGGC TCTGAGAGCT 50  
CTTGCAGTTG ATGGTGGCCC TCTCGCCC 78

## (2) INFORMATION FOR SEQ ID NO:58:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 30 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: unknown

77

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

GCGGAATTCG TAGTCGGATA TCGTGATGAC

30

(2) INFORMATION FOR SEQ ID NO:59:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 34 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:

TGGAAAGCTT GGCGCCGCCA CAGTACGTTT GATC

34

(2) INFORMATION FOR SEQ ID NO:60:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 57 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..57

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:60:

ATG GTG TTG CAG ACC CAG GTC TTC ATT TCT CTG TTG CTC TGG 42  
 Met Val Leu Gln Thr Gln Val Phe Ile Ser Leu Leu Leu Trp  
   1                  5                  10

ATC TCT GGT GCC TAC  
 Ile Ser Gly Ala Tyr  
   15

57

78

## (2) INFORMATION FOR SEQ ID NO:61:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 19 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:61:

Met	Val	Leu	Gln	Thr	Gln	Val	Phe	Ile	Ser	Leu	Leu	Leu	Trp	Ile
1				5					10					15
Ser Gly Ala Tyr														



## WHAT IS CLAIMED IS:

1. A fusion molecule comprising a first fusion partner nucleotide sequence encoding an amino acid sequence having the antigen specificity of a anti-*Plasmodium* antibody, operatively linked in frame to a second fusion partner nucleotide sequence.
2. The molecule according to claim 1 wherein said first fusion partner is a synthetic immunoglobulin variable region nucleotide sequence encoding an amino acid sequence comprising a complementarity determining region originating from a *Plasmodium* antibody, a fragment or allelic variation or modification thereof.
3. The molecule according to claim 2 wherein said second fusion partner is a heterologous immunoglobulin variable framework region.
4. The molecule according to claim 2 wherein said variable region nucleotide sequence is selected from the group consisting of a heavy chain variable region and a light chain variable region.
5. The molecule according to claim 4 selected from the group consisting of
  - (a) a heavy chain nucleotide sequence of Fig. 5 (SEQ ID NO: 11);
  - (b) a heavy chain nucleotide sequence of Fig. 6 (SEQ ID NO: 13);
  - (c) a light chain nucleotide sequence of Fig. 2 (SEQ ID NO: 5) and Fig. 3 (SEQ ID NO: 7).
  - (d) a light chain nucleotide sequence of Fig. 3 (SEQ ID NO: 7); and
  - (e) functional fragments thereof.

6. The molecule according to claim 4 wherein said first fusion partner nucleotide sequence comprises a sequence selected from the group consisting of

- (a) AGCTATGCCATGAGC: SEQ ID NO: 15;
- (b) GAAATTAGTGATGGTGGTAGTTACACCTACTATCCA  
GACACTGTGACGGGC: SEQ ID NO: 17;
- (c) CTCATCTACTATGGTTACGACGGGTATGCTATGGAC  
TAC: SEQ ID NO: 19;
- (d) AAGAGCTCTCAGAGCCTTTTATACTCGAGCAATCAA  
AAGAATTACTTGGCC: SEQ ID NO: 21;
- (e) TGGGCGTCAACTAGGGAATCT: SEQ ID NO: 23;
- (f) CAGCAATATTATAGCTATCCGCGGACG: SEQ ID  
NO: 25;
- (g) AGCTATGCCATGTCT: SEQ ID NO: 32;
- (h) AAGTCCAGTCAGAGCCTTTTATATAGTAGCAATCAAA  
AGAATTACTTGGCC: SEQ ID NO: 34;
- (i) TGGGCATCCACTAGGGAATCT: SEQ ID NO: 36;
- (j) CAGCAATATTATAGCTATCCTCGGACG: SEQ ID  
NO: 38;
- (k) TGGGCGTCGACTAGGGAATCT: SEQ ID NO: 41;

and an allelic variation or modification thereof, characterized by the antigen specificity of murine NFS2, said nucleic acid sequence optionally containing restriction sites to facilitate insertion into a desired antibody framework region or a plasmid vector.

7. A synthetic immunoglobulin variable region nucleotide sequence encoding an amino acid sequence comprising a complementarity determining region originating from a *Plasmodium* antibody, a fragment or allelic variation or modification thereof.

8. A fusion protein comprising a first amino acid sequence derived from a *Plasmodium* antibody capable of binding an epitope on a selected *Plasmodium* species, said sequence having the antigen specificity of said antibody fused to a heterologous second amino acid sequence.

9. The fusion protein according to claim 8 wherein said first amino acid sequence comprises an amino acid sequence selected from the group consisting of:

- (a) a variable heavy chain sequence of said antibody;
- (b) a variable light chain sequence of said antibody;
- (c) a complementarity determining region of said antibody; and
- (d) a functional fragment of (a) through (c).

10. The fusion protein according to claim 8 wherein said fusion protein is selected from the group consisting of

- (a) a complete engineered antibody, having full length heavy and light chains comprising at least fragments of the variable regions derived from said *Plasmodium* antibody;
  - (b) the  $F_{ab}$  or  $(F_{ab}')_2$  fragment of the engineered antibody of (a);
  - (c) a dimer formed of heavy chains derived from the engineered antibody of (a);
  - (d) an  $F_v$  fragment of the engineered antibody of (a); and
  - (e) a single-chain antibody derived from the engineered antibody of (a);
- said protein having the same specificity as said *Plasmodium* antibody.

11. An engineered *P. falciparum* antibody comprising a heavy chain comprising a complementarity determining region derived from the variable heavy chain region of a non-human *P. falciparum* monoclonal antibody.

12. The antibody according to claim 11 wherein said non-human CDRs are in operative association with one of the group consisting of

(a) a selected human antibody heavy chain framework and constant regions; and

(b) the heavy chain framework from said antibody and a constant region from a selected human antibody.

13. The antibody according to claim 11 further comprising a light chain selected from the group consisting of

(a) a light chain comprising a CDR derived from the variable light chain region of said monoclonal antibody in operative association with selected human antibody light chain framework and constant regions;

(b) the light chain framework from said antibody and a constant region from a selected human antibody;

(c) the complete light chain from said anti-*Plasmodium* antibody; and

(d) the complete light chain from a selected human antibody.

14. The antibody according to claim 11, wherein said heavy chain comprises a variable heavy chain sequence selected from the sequences of Fig. 5 (SEQ ID NO: 12), Fig. 6 (SEQ ID NO: 14), Pfzhzc2-3 (SEQ ID NO:14), and Pfzhzc2-6 (SEQ ID NO:42).

15. The antibody according to claim 13 wherein said light chain comprises a variable light chain sequence selected from the sequences of Fig. 2 (SEQ ID NO: 6) and Fig. 3 (SEQ ID NO: 8).

16. The antibody according to claim 13 wherein light chain complementarity determining region is selected from one or more of the sequences consisting of

- (a) LysSerSerGlnSerLeuLeuTyrSerSerAsn  
GlnLysAsnTyrLeuAla: SEQ ID NO: 22;
- (b) TrpAlaSerThrArgGluSer: SEQ ID NO: 24;  
and
- (c) GlnGlnTyrTyrSerTyrProArgThr: SEQ ID  
NO: 26.

17. The antibody according to claim 11 wherein said heavy chain complementarity determining region is selected from one or more of the sequences consisting of

- (a) SerTyrAlaMetSer: SEQ ID NO: 16;
- (b) GluIleSerAspGlyGlySerTyrThrTyrTyrPro  
AspThrValThrGly: SEQ ID NO: 18; and
- (c) LeuIleTyrTyrGlyTyrAspGlyTyrAlaMet  
AspTyr: SEQ ID NO: 20.

18. An anti-*P. falciparum* complementarity determining region peptide selected from the group consisting of

- (a) SerTyrAlaMetSer: SEQ ID NO: 16;
- (b) GluIleSerAspGlyGlySerTyrThrTyrTyr  
ProAspThrValThrGly: SEQ ID NO: 18;
- (c) LeuIleTyrTyrGlyTyrAspGlyTyrAlaMet  
AspTyr: SEQ ID NO: 20;
- (d) LysSerSerGlnSerLeuLeuTyrSerSerAsn  
GlnLysAsnTyrLeuAla: SEQ ID NO: 22;

(e) TrpAlaSerThrArgGluSer: SEQ ID NO: 24;

(f) GlnGlnTyrTyrSerTyrProArgThr: SEQ ID  
NO: 26;

and an analog thereof, characterized by the antigen  
specificity of NFS2.

19. A synthetic immunoglobulin variable chain  
amino acid sequence comprising a complementarity  
determining region originating from a *Plasmodium* antibody  
in a heterologous variable chain framework, a fragment or  
analog thereof sharing the anti-*Plasmodium* antigen  
specificity of said sequence.

20. The sequence according to claim 19  
selected from the group consisting of the amino acid  
sequences of Fig. 5 (SEQ ID NO: 12), Fig. 6 (SEQ ID NO:  
14), Fig. 2 (SEQ ID NO: 6) and Fig. 3 (SEQ ID NO: 8).

21. A monoclonal antibody, other than NFS2,  
which is capable of binding to a *P. falciparum* epitope  
comprising the sequence Pro Asn Ala Asn Pro Asn SEQ ID  
NO: 27, a F<sub>ab</sub> fragment thereof, or a (F<sub>ab</sub>')<sub>2</sub> fragment  
thereof.

22. A pharmaceutical prophylactic composition  
comprising a fusion protein or antibody according to any  
of claims 8 through 21 and a pharmaceutically acceptable  
carrier or diluent.

23. A pharmaceutical composition according to  
claim 22 wherein said protein is a humanized *P.*  
*falciparum* antibody.

24. A recombinant plasmid comprising a nucleic acid sequence of any of claims 1 through 7 in operative association with a regulatory control sequence capable of directing the replication and expression of said nucleic acid sequence in a selected host cell.

25. A mammalian cell line transfected with at least one recombinant plasmid comprising a nucleic acid sequence of any of claims 1 through 7.

26. A method of producing a engineered antibody comprising culturing a mammalian cell line transfected with at least one recombinant plasmid comprising a nucleic acid sequence of any of claims 1 through 7 under suitable conditions permitting expression and assembly of complementary heavy and light chains, and recovering the assembled antibody from the cell culture.

27. The use of a protein or antibody of claims 8 through 21 in the preparation of a pharmaceutical composition suitable for passively protecting a human against infection by a *Plasmodium* species.

1/9

FIGURE 1

[illegible]







4/9

FIGURE 4


CTC	GAG	TCT	GGG	GGA	GGC	TTA	GTG	AAG	CCT	30
Leu	Glu	Ser	Gly	Gly	Gly	Leu	Val	Lys	Pro	
1				5					10	
GGA	GGG	TCC	CTG	AAA	ATC	TCC	TGC	GCA	GCC	60
Gly	Gly	Ser	Leu	Lys	Ile	Ser	Cys	Ala	Ala	
				15					20	
TCT	GGA	TTC	ACT	TTC	AGT	AGC	TAT	GCC	ATG	90
Ser	Gly	Phe	Thr	Phe	Ser	<u>Ser</u>	<u>Tyr</u>	<u>Ala</u>	<u>Met</u>	
				25					30	
TCT	TGG	GTT	CGC	CAG	TCT	CCA	GAG	AAG	AGG	120
<u>Ser</u>	Trp	Val	Arg	Gln	Ser	Pro	Glu	Lys	Arg	
				35					40	
CTG	GAG	TGG	GTC	GCA	GAA	ATT	AGT	GAT	GGT	150
Leu	Glu	Trp	Val	Ala	<u>Glu</u>	<u>Ile</u>	<u>Ser</u>	<u>Asp</u>	<u>Gly</u>	
				45					50	
GGT	AGT	TAC	ACC	TAC	TAT	CCA	GAC	ACT	GTG	180
<u>Gly</u>	<u>Ser</u>	<u>Tyr</u>	<u>Thr</u>	<u>Tyr</u>	<u>Tyr</u>	<u>Pro</u>	<u>Asp</u>	<u>Thr</u>	<u>Val</u>	
				55					60	
ACG	GGC	CGA	TTC	ACC	ATC	TCC	AGA	GAC	AAT	210
<u>Thr</u>	<u>Gly</u>	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	
				65					70	
GCC	AAG	AAC	ACC	CTA	TAC	CTG	GAA	ATG	AGC	240
Ala	Lys	Asn	Thr	Leu	Tyr	Leu	Glu	Met	Ser	
				75					80	
AGT	CTG	AGG	TCT	GAG	GAC	ACG	GCC	ATG	TAT	270
Ser	Leu	Arg	Ser	Glu	Asp	Thr	Ala	Met	Tyr	
				85					90	
TAC	TGT	GCA	AGC	CTC	ATC	TAC	TAT	GGT	TAC	300
Tyr	Cys	Ala	Ser	<u>Leu</u>	<u>Ile</u>	<u>Tyr</u>	<u>Tyr</u>	<u>Gly</u>	<u>Tyr</u>	
				95					100	
GAC	GGG	TAT	GCT	ATG	GAC	TAC	TGG	GGT	CAA	330
<u>Asp</u>	<u>Gly</u>	<u>Tyr</u>	<u>Ala</u>	<u>Met</u>	<u>Asp</u>	<u>Tyr</u>	Trp	Gly	Gln	
				105					110	
GGA	ACC	TCA	GTC	ACC	GTC	TCC	TCA			354
Gly	Thr	Ser	Val	Thr	Val	Ser	Ser			
				115						

Figure 5

GAG	GTG	CAG	CTG	<u>Xho I</u>		TCT	GGG	GGA	GGC	TTG	GTA	CAG	39
Glu	Val	Gln	Leu	Leu	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	
1				5					10				
CCT	GGG	GGG	TCC	CTG	AGA	CTC	TCC	TGT	GCA	<u>Nhe I</u>		GGA	78
Pro	Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	
15						20					25		
TTC	ACC	TTT	AGC	AGC	TAT	GCC	ATG	AGC	TGG	GTC	CGC	CAG	117
Phe	Thr	Phe	Ser	<u>Ser</u>	<u>Tyr</u>	<u>Ala</u>	<u>Met</u>	<u>Ser</u>	Trp	Val	Arg	Gln	
			30					35					
GCT	CCA	GGG	AAA	<u>Xba I</u>		GAG	TGG	GTC	TCA	GAA	ATT	AGT	156
Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val	Ser	<u>Glu</u>	<u>Ile</u>	<u>Ser</u>	
40					45					50			
GAT	GGT	GGT	AGT	TAC	ACC	TAC	TAT	CCA	GAC	ACT	GTG	ACG	195
<u>Asp</u>	<u>Gly</u>	<u>Gly</u>	<u>Ser</u>	<u>Tyr</u>	<u>Thr</u>	<u>Tyr</u>	<u>Tyr</u>	<u>Pro</u>	<u>Asp</u>	<u>Thr</u>	<u>Val</u>	<u>Thr</u>	
		55					60					65	
GGC	CGG	TTC	ACG	<u>Eco RV</u>		AGA	GAC	AAT	TCC	AAG	AAC	ACG	234
<u>Gly</u>	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	
				70					75				
CTG	TAT	CTG	CAA	ATG	AAC	AGC	CTG	AGA	GCC	GAG	GAC	ACT	273
Leu	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	
	80					85					90		
<u>Pst I</u>													
GCA	GTA	TAT	TAC	TGT	GCG	AAA	CTC	ATC	TAC	TAT	GGT	TAC	312
Ala	Val	Tyr	Tyr	Cys	Ala	Lys	<u>Leu</u>	<u>Ile</u>	<u>Tyr</u>	<u>Tyr</u>	<u>Gly</u>	<u>Tyr</u>	
			95					100					
GAC	GGG	TAT	GCT	ATG	GAC	TAC	TGG	GGC	CAG	<u>Kpn I</u>		CTG	351
<u>Asp</u>	<u>Gly</u>	<u>Tyr</u>	<u>Ala</u>	<u>Met</u>	<u>Asp</u>	<u>Tyr</u>	Trp	Gly	Gln	Gly	Thr	Leu	
105					110					115			
<u>Sst I</u>													
GTC	ACC	GTG	AGC	TCA	GCTAGTACCA			AGGGCCCAAG			CTT		389
Val	Thr	Val	Ser	Ser									
		120											

Figure 6

GAG	GTG	CAG	CTG	<u>Xho I</u>		TCT	GGG	GGA	GGC	TTG	GTA	CAG	39
Glu	Val	Gln	Leu	Leu	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	
1				5					10				
CCT	GGG	GGG	TCC	CTG	AGA	CTC	TCC	TGT	GCA	<u>Nhe I</u>		GGA	78
Pro	Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	
15						20					25		
TTC	ACC	TTT	AGC	AGC	TAT	GCC	ATG	AGC	TGG	GTC	CGC	CAG	117
Phe	Thr	Phe	Ser	<u>Ser</u>	<u>Tyr</u>	<u>Ala</u>	<u>Met</u>	<u>Ser</u>	Trp	Val	Arg	Gln	
			30					35					
GCT	CCA	GGG	AAA	<u>Xba I</u>		TGG	GTC	TCA	GAA	ATT	AGT	156	
Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val	Ser	<u>Glu</u>	<u>Ile</u>	<u>Ser</u>	
40				45						50			
GAT	GGT	GGT	AGT	TAC	ACC	TAC	TAT	CCA	GAC	ACT	GTG	ACG	195
<u>Asp</u>	<u>Gly</u>	<u>Gly</u>	<u>Ser</u>	<u>Tyr</u>	<u>Thr</u>	<u>Tyr</u>	<u>Tyr</u>	<u>Pro</u>	<u>Asp</u>	<u>Thr</u>	<u>Val</u>	<u>Thr</u>	
			55				60					65	
GGC	CGG	TTC	ACG	ATA	TCC	AGA	GAC	AAT	TCC	AAG	AAC	ACG	234
<u>Gly</u>	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	
				70					75				
CTG	TAT	CTG	CAA	ATG	AAC	AGC	CTG	AGA	GCC	GAG	GAC	ACT	273
Leu	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	
	80					85					90		
<u>Pst I</u>													
GCA	GTG	TAT	TAC	TGT	GCA	TCT	CTC	ATC	TAC	TAT	GGT	TAC	312
Ala	Val	Tyr	Tyr	Cys	Ala	Ser	<u>Leu</u>	<u>Ile</u>	<u>Tyr</u>	<u>Tyr</u>	<u>Gly</u>	<u>Tyr</u>	
			95				100						
GAC	GGG	TAT	GCT	ATG	GAC	TAC	TGG	GGC	CAA	<u>Kpn I</u>		CTG	351
<u>Asp</u>	<u>Gly</u>	<u>Tyr</u>	<u>Ala</u>	<u>Met</u>	<u>Asp</u>	<u>Tyr</u>	Trp	Gly	Gln	GGT	ACC	Leu	
105					110					115			
<u>Sst I</u>													
GTC	ACC	GTG	AGC	TCA	GCTAGTACCA			AGGGCCCAAG			CTT	389	
Val	Thr	Val	Ser	Ser									
		120											






FIG. 8





## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US93/08435

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(5) : Please See Extra Sheet.

US CL : 435/240.2, 320.1, 240.27; 424/85.8; 530/387.1, 388.6.

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/240.2, 320.1, 240.27; 424/85.8; 530/387.1, 388.6.

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS, Biosis, Medline.

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP, O, 270,077 (Nakatani et. al.) 06 August 1988, see entire document.	6,16-18,25,26
X	The EMBO Journal, Volume 5, No. 7, issued 1986, Andrew J. Caton et. al., "Structural and functional implications of a restricted antibody response to a defined antigenic region on the influenza virus hemagglutinin", pages 1577-1587, see figures 4 and 5.	6, 16-18, 25,26

☒ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be part of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

16 December 1993

Date of mailing of the international search report

27 DEC 1993

 Name and mailing address of the ISA/US  
 Commissioner of Patents and Trademarks  
 Box PCT  
 Washington, D.C. 20231

Authorized officer

PAULA HUTZELL

Facsimile No. NOT APPLICABLE

Telephone No. (703) 308-0196

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US93/08435

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	The Journal of Immunology, Volume 139, No.7, issued 01 October 1987, Harout Dersimonian et. al, "Relationship of human variable region heavy chain germ-line genes to genes encoding anti-DNA autoantibodies", pages 2496-2501, see page 2498.	1-27
Y	Bull World Health Organ, 68 Suppl. issued 1990, Mellouk et. al. "Evaluation of an in vitro assay aimed at measuring protective antibodies against sporozoites", pages 52-59, see entire abstract.	1-27
Y	Proceedings of the National Academy of Sciences, Volume 86, issued December 1989, Queen et. al. "A humanized antibody that binds to the interleukin 2 receptor", pages 10029-10033, see entire document.	1-27

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US93/08435

## A. CLASSIFICATION OF SUBJECT MATTER:

IPC (5):

C07H 21/02, 21/04; C12N 15/70, 15/74, 15/79, 5/10; C07K 15/28; A61K 39/395.